

1 **RESEARCH ARTICLE**2 **Bioefficacy of Flupyradifurone 200 SL Against Jassids, Whiteflies and Their**
3 **Impact on Natural Enemies in Brinjal**

4 Sangamithra, S*, Vinothkumar, B. and Muthukrishnan, N

5 Department of Agricultural Entomology, Tamil Nadu Agricultural University, Coimbatore-641 003

6 Corresponding author agri.sangamithra@gmail.com

7
8**ABSTRACT**

Two field experiments were conducted at Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu to evaluate bioefficacy, phytotoxicity effect if any of flupyradifurone 200 SL against whiteflies and leaf hoppers and its safety to non target invertebrates in brinjal during 2015 – 16. The results revealed that flupyradifurone 200 SL@ 175 g ai ha⁻¹ was effective in reducing whiteflies and leaf hoppers population. The whiteflies per cent reduction of 96.40 and 95.77 was obtained in first and second season, respectively. Similarly, regarding leaf hopper 96.35 and 97.92 per cent reduction over control was recorded. The imposed treatments were found safer to non target invertebrates like spiders and coccinellids. Furthermore, no phytotoxic effect was observed even in treatments imposed with flupyradifurone 200 SL @ 75 and 150 g a.i. ha⁻¹.

9 **Keywords:** Flupyradifurone, whiteflies, leafhoppers, brinjal, safety.

10

11 **INTRODUCTION**

12 Brinjal (*Solanum melongena* Linn.) commonly known as egg plant and forever as “King of Vegetables”
13 (Thompson and Kelly, 1957) is cultivated extensively in India under diverse agro climatic conditions and
14 throughout the year (Nayak *et al.*, 2014). The production statistics of 2014-15 divulged that brinjal is
15 cultivated in 6, 80,000 ha with productivity of 12, 706, 000 tonnes (Saxena, 2015). It is ranked among top
16 ten vegetables in terms of antioxidant capacity and flavonoid constituents (Timberlake, 1981; Singh *et al.*,
17 2009) which have been associated with various health benefits (Ames *et al.*, 1993; Hung *et al.*, 2004). The
18 extracts from brinjal contain anthocyanins and strychnine which are effective in curing a number of
19 diseases including cancer, high blood pressure and hepatitis (Magioli and Mansur, 2005; Silva *et al.*,
20 1999). In view of ecological sustainability, brinjal is also not exempted from biotic stress and more than
21 30 insect pests are found to cause significant damage right from germination to harvest (Ragupathy *et al.*,
22 1997). Nevertheless shoot and fruit borer (*Leucinodes orbonalis* Gu.) is considered to be abnoxious,
23 sucking pests *viz.*, leaf hoppers (*Amrasca biguttula biguttula* Ishida and *A. devastans* Distant) and
24 whiteflies (*Bemisia tabaci* Gennadius) are under prime consideration that causes inflicting level of damage
25 and yield loss (Goshal and Chatterjee, 2013; Mahmood *et al.*, 2002; Kalawate and Dethe, 2012; Sultana *et al.*,
26 2012; Shrinivasan and Babu, 2001). Both leafhoppers and whiteflies are widely distributed in tropical,
27 sub tropical and temperate regions. The nymphs and adults suck sap from the lower surface of leaves, by
28 which nutrient translocation is disrupted in conducting vessels and also apparently introduce a toxin that
29 affects photosynthesis in proportion to the amount of feeding resulting in hopper burn. Besides, the
30 honeydew secretion attracts black sooty mold that impairs photosynthesis and moreover both pests are
31 considered to be potential vectors of copious viruses (Sharma and Chander, 1998).

32 The management of pests by insecticide application remains to be frontline and unsurpassed technique.
33 On the other hand as survival to the fittest, insects also develop resistance even for molecules that target
34 unique sites as mode of action. One such group of insecticides, recognized universally for the management
35 of sap sucking group of insects was neonicotinoids and copious numbers of insecticides had been
36 evaluated for efficacy against sucking pests alone and in combination with shoot and fruit borer. To cite
37 few efficacy findings in brinjal, imidacloprid @ 18 and 22.5 g ha⁻¹ and thiamethoxam @ 25 and 50 g ha⁻¹
38 against leafhopper and whitefly (Mhaske and Mote, 2005); Imidacloprid 70 WG @ 0.2 g L⁻¹, buprofezin 40
39 SC @ 2 mL L⁻¹ and fipronil 50 SC @ 2mL and 1 mL L⁻¹ against jassids and whiteflies (Das and Islam, 2014).
40 Imidacloprid 17.8 SL @ 0.5 mL L⁻¹ against aphids, leaf hoppers and whiteflies (Rajesh Kumar *et al.*, 2017);
41 Thiamethoxam @ 0.025%, diafenthiuron @ 0.05% , thiacloprid @ 0.012% against leaf hopper;
42 spiromesifen @ 0.024%, diafenthiuron @ 0.05% and triazophos @ 0.08% against whiteflies (Shaikh and
43 Patel, 2012); Flubendiamide 24% w/v + thiacloprid 24% sc w/v @ 84 + 84 g a.i ha⁻¹ against aphids,

44 jassids and thrips (Sangamithra *et al.*, 2018). Besides development of molecules with novel mode of
 45 action, crisis of resistance development is inevitable, hence standardization and commercialization of
 46 newer molecules of insecticides is mandate to attain sustainable pest management. New selective
 47 insecticides compatible with modern integrated pest management (IPM) principles addressing the
 48 regulatory needs for an improved toxicological and environmental profile will stepwise replace older
 49 chemistry suffering from resistance development in many invertebrate pests frequently targeted by
 50 indispensable chemical treatments in some agricultural settings (Nauen *et al.*, 2012). In search of new
 51 chemical scaffolds leading to novel chemical classes of insecticides, particularly for sucking pest control,
 52 the natural product stemofoline known as a potent agonist of insect nicotinic acetylcholine receptors
 53 (nAChRs), was considered as a good starting point and seed for the development of flupyradifurone, the
 54 first representative of the novel butenolide class of insecticides active against various sucking pests
 55 (Tamura *et al.*, 1978; Uvary, 1999). Its mode of action is similar to neonicotinoids that acts on nACh
 56 receptors, however chemical structure differs from nitroguanidine neonicotinoids, hence placed under sub
 57 group Group 4 in IRAC classification (Jeschke *et al.*, 2013; Jeschke and Nauen, 2008). In this regard, two
 58 field experiments were conducted during two consecutive years (2015 - 2016) to evaluate bioefficacy of
 59 flupyradifurone 200 SL, phytotoxicity if any and its impact to non - target organisms.

60 MATERIAL AND METHODS

61 Two field experiments were conducted at Tamil Nadu Agricultural University, Coimbatore during July -
 62 October 2015 and January - May 2016 (Variety: CO - 2). The experiment was laid out in Randomized
 63 Block Design with four replications. The treatment details were T1 - Flupyradifurone 200 SL @ 125 g a.i
 64 ha⁻¹; T2 - Flupyradifurone 200 SL @ 150 g a.i ha⁻¹; T3 - Flupyradifurone 200 SL @ 175 g a.i ha⁻¹;
 65 Phosphamidon 40 % SL @ 300 g a.i. ha⁻¹ and T5 - Untreated control. The treatments were imposed on 30
 66 days old crop and applied twice at weekly interval. The treatments were sprayed with pneumatic knapsack
 67 sprayer using 500 litres of spray fluid per hectare.

68 (i) Method of assessment of pest and natural enemies population

69 The population of sucking pests *viz.*, whiteflies (nymphs) and leaf hopper were recorded on six leaves per
 70 plant (three leaves at top and three leaves at bottom) of five randomly selected plants per plot prior to
 71 spraying followed by 3,7,10 and 15 days after each spray and expressed as number per six leaves. The day
 72 observations were pooled, mean population and per cent reduction over control was calculated after each
 73 spray. Brinjal yield per plot was recorded from each harvest and pooled to arrive at the total yield and
 74 expressed as tonnes ha⁻¹. Five randomly selected plants per plot were thoroughly observed for population
 75 of natural enemies. The population of the predators (coccinellids and spiders) was recorded before and 3,
 76 7, 10 and 15 days after each spraying and expressed as numbers per five plants.

77 (ii) Assessment of phytotoxicity

78 The plants were sprayed with flupyradifurone 200 SLw/w @ 175 and 350 g a.i. ha⁻¹ to assess the
 79 occurrence of phytotoxicity. The plants were observed on 1, 3, 7, 10, 14 and 21 days after spraying as per
 80 the protocol of Central Insecticide Board Registration Committee (C.I.B. and R.C). for the phytotoxic
 81 symptoms like injury to leaf tip and leaf surface, wilting, vein clearing, necrosis, epinasty and hyponasty
 82 which were recorded based on the following visual rating scale of 0 - 10 *viz.*, 0 - No phytotoxicity; 1 - 1-10
 83 %; 2 - 11-20 %; 3 - 21-30 %; 4 - 31-40 %; 5 - 41-50 %; 6 - 51-60 %; 7 - 61-70 %; 8 - 71-80 %; 9 - 81-90 %
 84 and 10 - 91-100 %. Per cent leaf injury was calculated using the formulae

$$\text{Per cent leaf injury} = \frac{\text{Total grade points}}{\text{Max. grade} \times \text{No. of leaves observed}} \times 100$$

85

86 (iii) Statistical analysis

87 The corrected per cent reduction of pest population over control in the field was worked out by using the
 88 formula given by Henderson and Tilton (1955).

$$\text{Corrected per cent reduction} = \left(1 - \left[\frac{T_a \times C_b}{T_b \times C_a} \right] \right) \times 100$$

89 where, T_a - Number of insects in the treatment after spraying; T_b - Number of insects in the treatment
 90 before spraying; C_b - Number of insects in the untreated check before spraying; C_a - Number of insects in
 91 the untreated check after spraying

92 The data on percentage was transformed into arc sine values and the population number into
 93 square root values before statistical analysis. The data obtained from field experiments were analysed in
 94 randomized block design (RBD) (Gomez and Gomez, 1984). The mean values were separated using
 95 Duncan's Multiple Range Test (DMRT) (Duncan, 1951).

96 **RESULTS AND DISCUSSION**

97 **(i) Bioefficacy of flupyradifurone 200 SL against whiteflies**

98 The population of whiteflies before application of treatments ranged from 15.70 to 16.35 six leaves⁻¹
99 (Table 1). Flupyradifurone 200 SL at 175 g a.i / ha significantly reduced whiteflies population and recorded
100 lowest mean population of 4.12 nymphs six leaves⁻¹ with 76.23 per cent reduction over control after first
101 spray. This was followed by flupyradifurone 200 SL at 150 g a.i / ha (5.19 nymphs six leaves⁻¹ with 70.03
102 per cent reduction over control) and flupyradifurone 200 SL at 125 g a.i / ha (6.46 nymphs six leaves⁻¹ with
103 62.70 per cent reduction over control). The standard check, phosphamidon 40% SL at 300 g a.i/ha also
104 reduced whitefly population to 6.68 nymphs six leaves⁻¹ with 61.43 per cent reduction over control . After
105 second application, similar trend in reduction was observed and flupyradifurone 200 SL at 175 g a.i / ha
106 reduced the population completely and recorded mean population of 0.78 nymphs six leaves⁻¹ followed by
107 flupyradifurone 200 SL at 150 g a.i / ha (1.73 nymphs six leaves⁻¹). The untreated check recorded the
108 whiteflies population of 21.54 nymphs six leaves⁻¹. Based on the per cent reduction in population over
109 untreated check, the order of efficacy of different insecticidal treatments were flupyradifurone 200 SL at
110 175 g a.i/ha (96.40%) > flupyradifurone 200 SL at 150 g a.i/ha (91.99%) > flupyradifurone 200 SL at 125
111 g a.i/ha (86.49%) > phosphamidon 40% SL at 300 g a.i/ha (85.78%)

112

113 During the second season experiment, the population of whiteflies before application of treatments ranged
114 from 17.00 to 18.33 six leaves⁻¹ (Table 1). Flupyradifurone 200 SL at 175 g a.i / ha significantly reduced
115 whiteflies population and recorded lowest mean population of 5.84 nymphs six leaves⁻¹ with 70.66 per
116 cent reduction over control after first spray. This was followed by flupyradifurone 200 SL at 150 g a.i / ha
117 (7.08 nymphs six leaves⁻¹ with 64.47 per cent reduction over control) and flupyradifurone 200 SL at 125 g
118 a.i / ha (8.25 nymphs six leaves⁻¹ with 54.56 per cent reduction over control). The standard check,
119 phosphamidon 40% SL at 300 g a.i/ha also reduced whitefly population to 8.44 nymphs six leaves⁻¹ with
120 57.63 per cent reduction over control. After second application, similar trend in reduction was observed
121 and flupyradifurone 200 SL at 175 g a.i / ha reduced the population completely and recorded mean
122 population of 1.01 nymphs six leaves⁻¹ followed by flupyradifurone 200 SL at 150 g a.i / ha (2.12 nymphs
123 six leaves⁻¹). The untreated check recorded the whiteflies population of 23.94 nymphs six leaves⁻¹. Based
124 on the per cent reduction in population over untreated check, the order of efficacy of different insecticidal
125 treatments were flupyradifurone 200 SL at 175 g a.i/ha (95.77%) > flupyradifurone 200 SL at 150 g a.i/ha
126 (91.15%) > flupyradifurone 200 SL at 125 g a.i/ha (84.98%) > phosphamidon 40% SL at 300 g a.i/ha
127 (83.35%)

128 **(ii) Bioefficacy of flupyradifurone 200 SL against leafhoppers**

129 The population of leafhoppers before application of treatments ranged from 13.25 to 14.00 six leaves⁻¹
130 (Table 2). Flupyradifurone 200 SL at 175 g a.i / ha significantly reduced leafhopper population and
131 recorded lowest mean population of 2.91 six leaves⁻¹ with 81.68 per cent reduction over control after first
132 spray. This was followed by flupyradifurone 200 SL at 150 g a.i / ha (4.26 six leaves⁻¹ with 73.20 per cent
133 reduction over control) and flupyradifurone 200 SL at 125 g a.i / ha (5.13 six leaves⁻¹ with 67.72 per cent
134 reduction over control). The standard check, phosphamidon 40% SL at 300 g a.i/ha also reduced
135 leafhopper population to 5.33 six leaves⁻¹ with 66.49 per cent reduction over control. After second
136 application, similar trend in reduction was observed and flupyradifurone 200 SL at 175 g a.i / ha reduced
137 the population completely and recorded mean population of 0.76 six leaves⁻¹ followed by flupyradifurone
138 200 SL at 150 g a.i / ha (1.91 six leaves⁻¹). The untreated check recorded the whiteflies population of
139 20.92 nymphs six leaves⁻¹. Based on the per cent reduction in population over untreated check, the order
140 of efficacy of different insecticidal treatments were flupyradifurone 200 SL at 175 g a.i/ha (96.35%) >
141 flupyradifurone 200 SL at 150 g a.i/ha (90.86%) > flupyradifurone 200 SL at 125 g a.i/ha (85.66%) >
142 phosphamidon 40% SL at 300 g a.i/ha (84.76%)

143

144 During the second season experiment, the population of leafhopper before application of treatments
145 ranged from 10.15 to 10.70 six leaves⁻¹ (Table 2). Flupyradifurone 200 SL at 175 g a.i / ha significantly
146 reduced leafhopper population and recorded lowest mean population of 3.31 six leaves⁻¹ with 77.12 per
147 cent reduction over control after first spray. This was followed by flupyradifurone 200 SL at 150 g a.i / ha
148 (4.50 six leaves⁻¹ with 68.83 per cent reduction over control) and flupyradifurone 200 SL at 125 g a.i / ha
149 (5.72 nymphs six leaves⁻¹ with 60.38 per cent reduction over control). The standard check, phosphamidon
150 40% SL at 300 g a.i/ha also reduced whitefly population to 6.18 six leaves⁻¹ with 57.22 per cent reduction
151 over control. After second application, similar trend in reduction was observed and flupyradifurone 200 SL
152 at 175 g a.i / ha reduced the population completely and recorded mean population of 0.43 six leaves⁻¹
153 followed by flupyradifurone 200 SL at 150 g a.i / ha (1.45 six leaves⁻¹). The untreated check recorded the
154 whiteflies population of 20.67 nymphs six leaves⁻¹. Based on the per cent reduction in population over

155 untreated check, the order of efficacy of different insecticidal treatments were flupyradifurone 200 SL at
156 175 g a.i./ha (97.92%) > flupyradifurone 200 SL at 150 g a.i./ha (93.00%) > flupyradifurone 200 SL at 125
157 g a.i./ha (86.81%) > phosphamidon 40% SL at 300 g a.i./ha (86.04%)

158 (iii) Impact on non target vertebrates and yield

159 The generalist predators that are commonly available in brinjal ecosystem viz., spiders and coccinellids
160 were chosen as non – target invertebrates and their population assessed to study the impact of
161 insecticide treatments. Flupyradifurone 200 SL, irrespective of doses found to be relatively safer to spiders
162 and coccinellids.

163

164 During first and season experiment the mean population after two rounds of spray indicated that
165 flupyradifurone 200 SL @ 125 g a.i. ha⁻¹ housed highest number of coccinellid population (5.43 and 5.30
166 five plants⁻¹, respectively) next to untreated check (8.03 and 8.65 five plants⁻¹, respectively). Subsequently
167 flupyradifurone 200 SL @ 150 g a.i. ha⁻¹ recorded 4.64 and 5.27 coccinellids five plants⁻¹, respectively. The
168 coccinellid population of 4.21 and 4.45 five plants⁻¹ observed in flupyradifurone 200 SL @ 175 g a.i. ha⁻¹.
169 The standard check phosphamidon 40% SL @ 300 g a.i. ha⁻¹ recorded coccinellid population of 4.11 and
170 4.42 five plants⁻¹ (Table 3). Regarding the spider population, comparable influence was exhibited by the
171 treatments in both the seasons. The mean population after two rounds of spray in both seasons revealed
172 that flupyradifurone 200 SL @ 125 g a.i. ha⁻¹ housed highest number of spider population (7.94 and 6.32
173 five plants⁻¹, respectively) next to untreated check (10.70 and 9.55 five plants⁻¹, respectively).
174 Flupyradifurone 200 SL @ 175 g a.i. ha⁻¹ recorded spider population as 6.66 and 4.95 five plants⁻¹,
175 respectively (Table 4).

176

177 The plants sprayed with flupyradifurone 200 SL @ 75 and 150 g a.i. ha did not show any phytotoxic
178 symptoms like leaf tip injury, wilting, vein clearing, necrosis, epinasty and hyponasty. The average fruit yield
179 in all the treatments ranged from 42.54 to 46.00 t ha⁻¹ during first season and 45.00 to 47.12 t ha⁻¹
180 during second season whereas 41.60 and 42.38 t ha⁻¹, respectively was observed in untreated control in
181 both seasons. Among, treatment imposed with flupyradifurone 200 SL @ 175 g a.i ha⁻¹ recorded highest
182 yield of 46.00 and 47.12 t ha⁻¹, respectively in two consecutive seasons (Table 5).


183

184 The contemporary experimental results infers flupyradifurone 200 SL @ 175 g a.i ha⁻¹ followed by 150 g
185 a.i. ha⁻¹ as effective dose for the management of leaf hopper and white fly population in brinjal. Together
186 with greater efficacy in pest management, it exhibits good safety profile for generalist predators viz.,
187 spiders and coccinellids in brinjal ecosystem. The precedent findings regarding flupyradifurone in brinjal
188 and other crops also depict the same conclusions which are discussed hereunder. The efficacy of
189 flupyradifurone 200 SL against rosy apple aphid (*Dysaphis plantaginea* (Passerini)) and green apple apid
190 (*Aphis pomi* (De Geer)) was investigated (Alston and Lindstrom, 2012). Flupyradifurone 200 SL @ 5.2 and
191 8.7 oz per acre performed well in reducing *D. plantaginea* faster and efficacy persisted for longer time (26
192 days post treatment). None of the treatments significantly reduced predator densities and recorded mean
193 as 1.2 total predators per shoot. Parasitism was significantly reduced in all insecticide treatments as
194 compared to the untreated control but as per authors, this was primarily caused by the higher densities of
195 rosy apple aphids available in the untreated control plots. Flupyradifurone 200 SL possess waiting period
196 of 15 days as similar to imidacloprid, while in management of mulberry thrips and toxicants does not show
197 any deleterious effects on growth of silkworm larvae as evidenced through non-significant differences in
198 economic and survival parameters of mulberry silkworm (Patil *et al.*, 2013). Flupyradifurone 20 SC @ 200
199 g a.i ha⁻¹ was elucidated as an effective alternate to neonicotinoids in cotton ecosystem (Rao *et al.*, 2014).
200 Flupyradifurone 200 SL @ 250 and 200 g a.i. ha⁻¹ provided superior control against leaf hoppers, aphids,
201 whiteflies and the population reduction was finer than neonicotinoids viz., imidacloprid 200 SL @ 20g
202 ai/ha and acetamiprid 20 SP @ 20 g a.i. ha⁻¹ even at lower dose of 150 g a.i ha⁻¹. Besides their efficacy,
203 highest seed cotton yield was obtained and did not influence population of natural enemies (Prasad,
204 2017). Flupyradifurone 200 SL @ 125, 150 and 175 g a.i ha⁻¹ was reported as effective management
205 practice for leafhoppers and whiteflies against standard, phosphamidon 40% SL @ 300 g a.i.
206 ha⁻¹ in brinjal at Rahuri, Maharashtra. The highest yield of brinjal fruits i.e. 76.96 and 79.03 q
207 ha⁻¹, respectively was recorded in flupyradifurone 200 SL @ 150 and 175 g a.i. ha⁻¹ and also found
208 relatively safer to coccinellid population in brinjal ecosystem (Wale *et al.*, 2017). Similar findings were
209 reported in brinjal at Vidisha, Madhya Pradesh (Vinod Kumar Garg *et al.*, 2018).


210

211

212 In the above field investigations of flupyradifurone 200 SL, the mainstream to be observed is use of
 213 neonicotinoids as comparable standard check and superiority of flupyradifurone. Nevertheless both groups
 214 of compounds seems to have same mode of action as an nAChR agonist, flupyradifurone is depicted as an
 215 effective alternate tool in resistance management strategies especially to sucking pest species that
 216 developed resistance to virtually all chemical classes of insecticides introduced to control them (Bass et
 217 al., 2014). In this regard, higher efficacy may be attributed towards its unique structural moiety known as
 218 butenolide that had been developed from natural product stemofoline. Stemofoline, isolated from leaves
 219 and stem of oriental medicinal plant *Stemona japonica* (Blume) Miq. (Stemonaceae) shows fast-acting
 220 insecticidal, antifeedant and repellent activities, but its activity is significantly lower than that of
 221 commercial products acting on insect nAChRs (Kaltenegger et al., 2003; Jeschke et al., 2013;
 222 Mungkornasawakul et al., 2004). Therefore, stemofoline was broadly used as a potent lead structure for
 223 development of novel active ingredients like flupyradifurone.



224 Fig 1. Natural product stemofoline 1 as the lead structure for novel ligands
 225
 226



227 Fig 2. New bioactive scaffold versus pharmacophore systems of known nAChR agonists
 228 (Nauen et al., 2015)
 229
 230

231 In the above fig. 1 head group which was identified as butenolide scaffold undergoes certain chemical
 232 changes to form enamino-carbonyl compound. It undergoes further chemical evolution via the butenolide
 233 subclasses resulting in discovery of flupyradifurone (Nauen et al., 2015).

234
 235 Furthermore distinct moiety of flupyradifurone can be explained by comparing with already commercialised
 236 nAChR agonists such as N-cyanoamidines (acetamiprid, thiacloprid), nitroenamines (nitenpyram), N-
 237 nitroguanidines (imidacloprid, clothianidin, thiamethoxam or dinotefuran) or sulfoximines (sulfoxaflor), the
 238 butenolide flupyradifurone 4 (Z =O) contains a different pharmacophore system as a new bioactive scaffold
 239 (Fig. 2). Besides, distinct structural moiety, efficacy of flupyradifurone may be explored with their agonist
 240 affinity and relative efficacy. Radioligand [³H] imidacloprid displacement studies was conducted to depict
 241 binding site and affinity of flupyradifurone and efficacy was inferred by whole cell clamp technique (Patch
 242 clamping) with holding potential of -70 mV. Results inferred that flupyradifurone displaces [³H] imidacloprid
 243 bound to *Musca domestica* (Linn.) nAChRs from its binding site with nanomolar affinity, and an I₅₀ value of
 244 2.38 ±1.93 nM was calculated. It activates endogenously expressed insect nAChRs by reverse binding and
 245 acts as a partial agonist with a relative agonist efficacy of 0.56 relative to the amplitude elicited by 1 mM
 246 of acetylcholine (Nauen et al., 2015). Flupyradifurone shows good translocation in short time after

247 application in planta, hence suggesting a good systemic activity. It is mainly translocated in the xylem, as
248 shown by its accumulation in distal leaf regions when taken up by the leaf lamina, roots and stems. Rapid
249 action on sucking pests, was exemplified by a translaminar study on the suppression of honeydew
250 excretion in green peach aphid feeding on the abaxial site of adaxially treated oilseed rape leaves (Nauen
251 *et al.*, 2015). Within a short time interval, most of the aphids stopped feeding and died 2 days later,
252 suggesting a high potential of flupyradifurone to prevent the transmission of plant pathogenic viruses at
253 recommended field rates. It can be foliarly applied even during flowering, as it shows no adverse effects on
254 actively foraging honey bees in long-term field trials in oilseed rape when applied at rates as high as 205 g
255 ha⁻¹ (Nauen *et al.*, 2015)

256

257 The distinct chemical structure of the novel butenolide pharmacophore and the lack of metabolic cross-
258 resistance of flupyradifurone led to the formation of a new subgroup (4D) within the IRAC mode-of-action
259 classification and evolve as a tool for setting up resistance management strategies based on mode of
260 action. Hence, flupyradifurone, occupies a place of best alternative to neonicotinoids and can be
261 recommended for management of sucking pests in brinjal.

262 CONCLUSION

263 Flupyradifurone @ 150 and 175 g a.i. ha⁻¹ exhibit the excellent control of sucking pests in brinjal
264 agroecosystem without causing any phytotoxicity to the plant. The distinct chemical structure of the novel
265 butenolide pharmacophore and the lack of metabolic cross-resistance of flupyradifurone led to the
266 formation of a new subgroup (4D) within the IRAC mode of action classification and evolve as a tool for
267 setting up resistance management strategies based on mode of action. Hence, flupyradifurone, occupies a
268 place of best alternative to neonicotinoids and can be recommended for management of sucking pests in
269 brinjal.

270 REFERENCES

- 271 Falston, D. and T. Lindstrom. 2012. Rossy apple aphid insecticide efficacy trial. Report – Apple Aphid
272 insecticide, Utah State University, p 26 -27.
- 273 Ames, B.N., Shigenaga, M.K. and T. M. Hagen. 1993. Oxidants, antioxidants and the degenerative
274 diseases of aging. *Proc. Natl. Acad. Sci.*, **90**: 7915 – 7922.
- 275 Bass, C., Puinean, A.M., Zimmer, C.T., Denholm, I., Field, L.M. and S. P. Foster. 2014. The evolution of
276 insecticide resistance in the peach potato aphid, *Myzus persicae*. *Insect Biochem Mol Biol.*, **51**: 41
277 – 51.
- 278 Das, G and T. Islam. 2014. Relative efficacy of some newer insecticides on the mortality of jassid and
279 whitefly in brinjal. *Int J Res Biol Sci.*, **4(3)**: 89 – 93.
- 280 Duncan, D.B. 1951. A significance test for differences between ranked treatment means in an analysis of
281 variance. *Va. J. Sci.*, **2**: 171-189.
- 282 Gomez, K.A. and A. A. Gomez. 1984. Statistical procedures for agricultural research. Wiley International
283 Science Publications. John Wiley and Sons, New York, pp.207-215.
- 284 Goshal, A. and M. L. Chatterjee. 2013. Evaluation of imidacloprid 17.8 SL against brinjal jassids. *J Entomol*
285 *Res.*, **37 (4)**: 319 – 323.
- 286 Henderson, C.F. and E. N. Tilton 1955. Tests with acaricides against the brown wheat mite. *J. Econ.*
287 *Entomol.*, 48: 157-161.
- 288 Hung, H.C., Josphipura, K.J., Jiang, R., Hu, F.B., Hunter, D. and S. A. Smith – Warner. 2004. Fruit and
289 vegetable intake and risk of major chronic disease. *J. Nat. Cancer Inst.*, **96**: 1577 – 1584.
- 290 Jeschke, P. and R. Nauen. 2008. Review: neonicotinoids – from zero to hero in insecticide chemistry. *Pest*
291 *Manag Sci.*, **64**: 1084 – 1098.
- 292 Jeschke, P., Nauen, R. and M.E. Beck. 2013. Nicotinic acetylcholine receptor agonists; a milestone for
293 modern crop protection. *Angew Chem In ted.*, **52**: 9464 – 9485.
- 294 Kalawate, A. and M. D. Deth. 2012. Bioefficacy study of biorational insecticides on brinjal. *J Biopesticides.*
295 **5(1)**: 75 - 80.
- 296 Kaltenecker, E., Brem, B., Mereiter, K., Kalchauer, H., Kahling, H. and O. Hofer. 2003. Insecticidal pyridol
297 [1,2 - a] azepine alkaloids and related derivatives from *Stemona* species. *Phytochemistry.* **77**:
298 803 – 816.
- 299 Magioli, C. and E. Mansur. 2005. Eggplant (*Solanum melongena* L.): Tissue culture, genetic transformation
300 and use as an alternative model plant. *Acta bot. bras.*, **19(1)**: 139 – 148.

- 301 Mahmood, T., Hussain, S.I., Khokhar, K.M., Jeelani, G. and Ahmed Mukhtar. 2002. Population dynamics of
302 leafhopper (*Amrasca biguttula biguttula*) on brinjal and effects of abiotic factors on it. *Asian J*
303 *Plant Sci.* **1(4)**: 403 - 404.
- 304 Mhaske, B.M. and U. N. Mote. 2005. Studies on evaluation of new insecticides against brinjal pest
305 complex. *J. Maharashtra Agri. Univ.*, **30(3)**: 303-306.
- 306 Mungkornasawakul, P., Pyne, S.G., Jatisatiener, A., Supyen, D., Jatisatiener Ch. And Lie We. 2004.
307 Phytochemical and larvicidal studies on *Stemona curtisii*: structure of a new pyrido [1,2 - a]
308 azepine *Stemona* alkaloid. *J Nat Prod.*, **67**: 675 – 677.
- 309 Nauen, R., Elber, A., McCaffery, A.R., Slater, R. and T.C. Sparks. 2012. IRAC: insecticide resistance and
310 mode of action classification of insecticides in modern crop protection compounds. In:
311 Insecticides, W Kramer, U Schirmer, P Jeschke and M Witschel (Eds), Wiley – VCH, Weinheim,
312 Germany, Vol 3, 2nd Edn, pp 935 – 955.
- 313 Nauen, R., Jeschke, P., Robert, V., Beck, M.E., Ebbinghaus – Kintscher, U., Thielert, W., Wolfel, K., Haas, M.,
314 Kunz, K. and G. Raupach. 2015. Flupyradifurone: a brief profile of a new butenolide insecticide.
315 *Pest Manage Sci.* **71**: 850 – 862.
- 316 Nayak, U.S., Baral, K., Mandal, P. and S. Chatterjee. 2014. Influence of environmental factors on
317 population dynamics and infestation pattern of *Leucinodes orbonalis* in winter brinjal of Odisha. *J*
318 *Bioresource Stress Manage* **5 (3)**: 409 – 412.
- 319 Patil, Jyothi, Ashoka, J., Bheemanna, M., Nagangouda, A., Sreenivas, A.G. and Jaysree Mekali. 2013.
320 Waiting period for insecticide and a botanical used in control of mulberry thrips. *Annals Pl Pro.*
321 *Sci.*, **21**: 42 -45.
- 322 Prasad, N.V.V.S.D. 2017. Bioefficacy of novel insecticide flupyradifurone 200 SL against leaf hoppers,
323 aphids and whitefly in cotton. In: 19th Int Conf on Entomol, Paris, France on 19-20, 2017.
- 324 Ragupaty, A., Palanisamy, S., Chandramohan, N. and K. Gunathilagaraj. 1997. A guide on crop pests,
325 Sooriya Desk Top, Coimbatore, p 264.
- 326 Rajesh Kumar, Mahla, M.K., Beerendra Singh, Ahir, K.C. and N. C. Rathor. 2017. Relative efficacy of newer
327 insecticides against sucking insect pests of brinjal (*Solanum melongena*). *J Entomol Zool Studies.*,
328 **5(4)**: 914 – 917.
- 329 Rao, G.M.V., Prasad, N.V.V.S.D., Prasad, M., Malyadri and V. Chenga Reddy. 2014. Emerging trends in
330 insect pest scenario of Bt cotton and ecofriendly approaches for management. In: Nat. Sym. On
331 Emerging trends in Eco – friendly Insect pest Management, Department of Agricultural
332 Entomology, TNAU, Coimbatore, p 216 -217.
- 333 Sangamithra, S., Vinothkumar, B., Muthukrishnan, N. and T. Manoharan. 2018. Evaluation of bioefficacy of
334 flubendiamide 24% w/v + thiacloprid 24% SC w/v against shoot and fruit borer and its sucking
335 pests and its safety to non target organisms in brinjal. *J Entomol Zool Studies.*, **6(1)**: 245 – 249.
- 336 Saxena, M. 2015. *Horticulture statistics at a glance*, Oxford University Press, New Delhi.
- 337 Shaikh, A.A. and J. J. Patel. 2012. Bio – efficacy of insecticides against sucking pests in brinjal. *AGRES – An*
338 *International e .Journal* (ISSN 2277-9663) **1(4)**: 423 – 434.
- 339 Sharma, K. and S. Chander. 1998. Spatial distribution of jassid *Amrasca biguttula biguttula* (Ishida) on
340 cotton. *Indian J Ent.*, **60(4)**: 326 – 328.
- 341 Shrinivasan, G and P. C. S. Babu. 2001. Field evaluation of neem products against whitefly, *Bemisia tabaci*
342 *Gennadius* on brinjal. *Annals Pl Pro. Sci.*, **9**: 19 - 21.
- 343 Silva, M.E., Santos, R.C., O' Leary, M.C. and R. S. Santos. 1999. Effect of aubergine (*Solanum melongena*)
344 on serum and hepatitis cholesterol and triglycerides in rats. *Braz. Arch. Biol. Technol.*, **42**: 339 –
345 342.
- 346 Singh, A.P., Luthria, D., Wilson, T., Vorsa, N., Singh, V., Banuelos, G.S. and S. Pasakdee. 2009. Polyphenols
347 content and antioxidant capacity of eggplant pulp. *Food Chem.*, **114**: 955 – 961.
- 348 Sultana, R., Solangi, B.K., Bughio, B.A. and M. S. Wagon. 2012. Field evaluation of bio pesticides against
349 jassids on brinjal crop. *Sindhu Univ Res J.* **44(3)**: 439 – 440.
- 350 Tamura, S., Sakata, K. and A. Sakurai. 1978. Stemofoline as insecticide. Stemofoline as insecticide. JP
351 53127825 A, *Institute of physical and Chemical Research*, Japan.
- 352 Thompson, C.H. and C. W. Kelly. 1957. *Vegetable Crops*, McGraw Hill Book Co. Inc, USA, p 501.
- 353 Timberlake, C.F. 1981. Anthocyanins in Fruit and Vegetables. In: *Recent Advances in the Biochemistry of*
354 *Fruit and Vegetables*, J Friend and M. J. C. Rhodes (Eds), Academic Press, New York, USA, pp 221-
355 47.

356 Uvary, I. 1999. Nicotine and other insecticidal alkaloids. In: Neonicotinoid Insecticides and the Nicotinic
 357 Acetylcholine Receptor, I Yamamoto and JE casida (Eds), Springer – Verlag, Berlin, Heidelberg,
 358 Newyork, pp 29-69.

359 Vinod Kumar Garg, Yogesh Patel, Raghuwanshi, M.S. and G. S. Jamliya. 2018. Bioefficacy of
 360 flupyradifurone 200 SL against sucking pest and their natural enemies on brinjal (*Solanum*
 361 *melongena* L.). *Annals of PI Soil Res.*, **20(1)**: 73 -76.

362 Wale, S.G., Pawar, S.A. and R.V. Datkhile . 2017. Evaluation of flupyradifurone 200 SL against sucking
 363 pests on Brinjal. *Annals PI Pro. Sci.*, **25** (2): 254 – 258.

364

365 **Table 1. Effect of flupyradifurone 200 SL against whitefly in brinjal**

366

Treatments	Number of whitefly nymphs six leaves ⁻¹											
	I Season						II Season					
	First spray			Second spray			First spray			Second spray		
	PTC	Mean*	PRC	PTC	Mean*	PRC	PTC	Mean*	PRC	PTC	Mean*	PRC
Flupyradifurone 200 SL @ 125 g a.i./ha	15.80	6.46 (2.54) ^c	62.70	6.28	2.91 (1.71) ^c	86.49	17.00	8.25 (2.87) ^c	58.56	7.10	3.60 (1.90) ^c	84.98
Flupyradifurone 200 SL @ 150 g a.i./ha	16.20	5.19 (2.28) ^b	70.03	5.15	1.73 (1.32) ^b	91.99	17.12	7.08 (2.66) ^b	64.47	5.80	2.12 (1.46) ^b	91.15
Flupyradifurone 200 SL @ 175 g a.i./ha	15.90	4.12 (2.03) ^a	76.23	4.07	0.78 (0.88) ^a	96.40	18.04	5.84 (2.42) ^a	70.66	4.58	1.01 (1.00) ^a	95.77
Phosphamidon 40% SL @ 300 g a.i./ha	16.35	6.68 (2.58) ^d	61.43	6.40	3.06 (1.75) ^d	85.78	17.50	8.44 (2.91) ^d	57.63	7.32	3.99 (2.00) ^d	83.35
Untreated check	15.70	17.33 (4.16) ^e	-	18.95	21.54 (4.64) ^e	-	18.33	19.92 (4.46) ^e	-	21.06	23.94 (4.89) ^e	-
SE. d	-	0.01	-	-	0.02	-	-	0.01	-	-	0.03	-
CD (P = 0.05)	-	0.03	-	-	0.04	-	-	0.03	-	-	0.07	-

367 *Mean of four observations; Values in parantheses are square root transformed values; In a column, means followed
 368 by a common letter are not significantly different by LSD (P=0.05) PTC- Pretreatment count; DAT – Days after
 369 treatment; PRC – Percent reduction over control

370

371

372 **Table 2. Effect of flupyradifurone 200 SL against leaf hoppers in brinjal**

373

Treatments	Number of leaf hoppers six leaves ⁻¹											
	I Season						II Season					
	First spray			Second spray			First spray			Second spray		
	PTC	Mean*	PRC	PTC	Mean*	PRC	PTC	Mean*	PRC	PTC	Mean*	PRC
Flupyradifurone 200 SL @ 125 g a.i./ha	13.95	5.13 (2.26) ^c	67.72	5.00	3.00 (1.73) ^c	85.66	10.15	5.72 (2.39) ^c	60.38	4.56	2.73 (1.65) ^c	86.81
Flupyradifurone 200 SL @ 150 g a.i./ha	14.00	4.26 (2.06) ^b	73.20	4.56	1.91 (1.38) ^b	90.86	10.58	4.50 (2.12) ^b	68.83	3.48	1.45 (1.20) ^b	93.00
Flupyradifurone 200 SL @ 175 g a.i./ha	13.70	2.91 (1.71) ^a	81.68	3.20	0.76 (0.87) ^a	96.35	10.34	3.31 (1.82) ^a	77.12	2.60	0.43 (0.66) ^a	97.92
Phosphamidon 40% SL @ 300 g a.i./ha	13.50	5.33 (2.31) ^d	66.49	5.12	3.19 (1.79) ^d	84.76	10.46	6.18 (2.49) ^d	57.22	5.00	2.89 (1.70) ^d	86.04
Untreated check	13.25	15.90 (3.99) ^e	-	18.00	20.92 (4.57) ^e	-	10.70	14.45 (3.80) ^e	-	17.20	20.67 (4.55) ^e	-
SE. d	-	0.01	-	-	0.01	-	-	0.02	-	-	0.01	-
CD (P = 0.05)	-	0.02	-	-	0.03	-	-	0.04	-	-	0.02	-

374 *Mean of four observations; Values in parantheses are square root transformed values; In a column, means followed
 375 by a common letter are not significantly different by LSD(P=0.05) PTC- Pretreatment count; DAT – Days after treatment;
 376 PRC – Percent reduction over control

377

378
379

Table 3. Safety of flupyradifurone 200 SL to coccinellids in brinjal ecosystem

Treatments	Number of coccinellids five plants ⁻¹											
	I Season						II Season					
	First spray			Second spray			First spray			Second spray		
	PTC	Mean*	PRC	PTC	Mean*	PRC	PTC	Mean*	PRC	PTC	Mean*	PRC
Flupyradifurone 200 SL @ 125 g a.i./ha	6.35	5.88 (2.42) ^b	14.16	6.40	5.43 (2.33) ^b	32.38	7.08	6.16 (2.48) ^b	20.72	6.48	5.30 (2.30) ^b	38.73
Flupyradifurone 200 SL @ 150 g a.i./ha	6.55	4.89 (2.21) ^c	28.61	5.25	4.64 (2.15) ^c	42.22	7.14	6.07 (2.46) ^b	21.88	6.40	5.27 (2.30) ^b	39.08
Flupyradifurone 200 SL @ 175 g a.i./ha	6.25	4.30 (2.07) ^d	37.23	4.85	4.21 (2.05) ^d	47.57	7.00	4.99 (2.23) ^c	35.78	5.20	4.45 (2.11) ^c	48.55
Phosphamidon 40% SL @ 300 g a.i./ha	6.50	4.08 (2.02) ^e	40.44	4.25	4.11 (2.03) ^d	48.82	7.46	4.91 (2.22) ^c	36.81	5.18	4.42 (2.10) ^c	48.90
Untreated check	5.95	6.85 (2.62) ^a	-	7.45	8.03 (2.83) ^a	-	7.39	7.77 (2.79) ^a	-	8.26	8.65 (2.94) ^a	-
SE. d	-	0.01	-	-	0.02	-	-	0.01	-	-	0.03	-
CD (P = 0.05)	-	0.02	-	-	0.05	-	-	0.02	-	-	0.06	-

380 *Mean of four observations; Values in parantheses are square root transformed values; In a column, means followed
381 by a common letter are not significantly different by LSD (P=0.05) PTC- Pretreatment count; DAT - Days after
382 treatment; PRC - Percent reduction over control

383
384

Table 4. Safety of flupyradifurone 200 SL to spiders in brinjal ecosystem

385
386

Treatments	Number of spiders five plants ⁻¹											
	I Season						II Season					
	First spray			Second spray			First spray			Second spray		
	PTC	Mean*	PRC	PTC	Mean*	PRC	PTC	Mean*	PRC	PTC	Mean*	PRC
Flupyradifurone 200 SL @ 125 g a.i./ha	9.55	8.50 (2.92) ^b	14.40	9.05	7.94 (2.82) ^b	25.79	8.04	6.98 (2.64) ^b	18.08	7.34	6.32 (2.51) ^b	33.82
Flupyradifurone 200 SL @ 150 g a.i./ha	9.40	8.10 (2.85) ^c	18.43	8.4	7.26 (2.69) ^c	32.15	8.33	6.75 (2.60) ^b	20.77	7.00	6.15 (2.48) ^b	35.60
Flupyradifurone 200 SL @ 175 g a.i./ha	8.55	7.26 (2.69) ^d	26.89	7.7	6.66 (2.58) ^d	37.76	8.56	6.03 (2.46) ^c	29.23	6.28	4.95 (2.22) ^c	48.17
Phosphamidon 40% SL @ 300 g a.i./ha	9.80	6.80 (2.61) ^e	31.52	6.95	6.25 (2.50) ^e	41.59	8.11	5.76 (2.40) ^c	32.39	5.80	4.87 (2.21) ^c	49.01
Untreated check	9.30	9.93 (3.15) ^a	-	10.25	10.70 (3.27) ^a	-	8.00	8.52 (2.92) ^a	-	9.06	9.55 (3.09) ^a	-
SE. d	-	0.03	-	-	0.02	-	-	0.03	-	-	0.02	-
CD (P = 0.05)	-	0.06	-	-	0.04	-	-	0.06	-	-	0.04	-

387 *Mean of four observations; Values in parantheses are square root transformed values; In a column, means followed
388 by a common letter are not significantly different by LSD(P=0.05) PTC- Pretreatment count; DAT - Days after treatment;
389 PRC - Percent reduction over control

390

391 **Table 5. Effect of flupyradifurone 200 SL on fruit yield in brinjal**
 392

Treatments			Yield* (t ha ⁻¹)	
			I Season	II Season
Flupyradifurone @ 125 g a.i./ha	200	SL	43.28 ^b	45.40 ^b
Flupyradifurone @ 150 g a.i./ha	200	SL	45.70 ^a	47.00 ^a
Flupyradifurone @ 175 g a.i./ha	200	SL	46.00 ^a	47.12 ^a
Phosphamidon @ 300 g a.i./ha	40%	SL	42.54 ^c	45.00 ^c
Untreated check			41.60 ^d	42.38 ^d
SE(d)			0.38	0.41
CD (P = 0.05)			0.76	0.82

393 *Mean of four observations; In a column, means followed by a common letter are not significantly
 394 different by LSD(P=0.05)
 395