



RESEARCH ARTICLE

Evaluation of efficient transformation method for *xylose reductase* gene integration in *Pichia pastoris* GS115

Yamunasri P¹, Priyadharshini R¹ and Sivakumar Uthandi*

¹Biocatalysts laboratory, Department of Agricultural Microbiology, Directorate of Natural Resource Management, Tamil Nadu Agricultural University, Coimbatore -641 003

ABSTRACT

Pichia pastoris remains an amicable host for the expression of heterologous protein from eukaryotes. Achieving a high level of expression of yeast in the recombinant host has remained a challenge. Apart from the codon preference of genes and proper protein folding for efficient expression of the heterologous gene, the main bottleneck is to efficiently integrate the heterologous gene into the host DNA. The present study evaluated the choice of available transformation methods such as lithium chloride, electroporation, and spheroplast for efficient integration of *xylose reductase* (*Xr*) gene of *Candida tropicalis* in *P. pastoris*. The results revealed that electrotransformation was found to be an efficient method for the transformation for *xr* in *P. pastoris*, yielding 1×10^6 transformants, followed by the lithium chloride method registering 2×10^3 transformants.

Received : 02nd November, 2020

Revised : 26th November, 2020

Revised : 08th December, 2020

Accepted : 24th December, 2020

Keywords: *Pichia pastoris*; *xylose reductase*; Transformation.

INTRODUCTION

D-Xylose is metabolized in yeast via., oxido-reductase pathway requiring several enzymes and cofactors. Xylose reductase (XR) is the first enzyme that converts D-xylose to xylitol requiring NAD(P)H, followed by the conversion of D-xylitol to D-xylulose by xylitol dehydrogenase (XDH) and requires NAD(P)⁺. Xylitol kinase then converts D-xylulose to D-xylulose-5-phosphate, which enters the pentose phosphate pathway (Jung *et al.*, 2015). XR and XDH are critical enzymes in the pathway. Xylitol is an economically important compound and has many applications in the food and pharmaceutical industry (Mohamad *et al.*, 2015). Since xylitol production from yeast is less expensive than chemical methods, many studies focused on the biological production of xylitol. Hence, cloning and integration of XR in yeast is vital in the production of xylitol. Introducing exogenous DNA (transformation) in yeast *Saccharomyces cerevisiae* was developed in 1978. Since then, many yeast and fungi have been successfully transformed with foreign DNA. There are chemical, physical, or enzymatic treatment methods for yeast transformation. A chemical methods employing lithium acetate or lithium chloride and PEG are well-known (Ito *et al.*, 1983). Similarly, the electroporation method of transformation is the preferred physical method when compared to the chemical and biolistic, and glass bead method. Enzyme-based methods generate spheroplasts, which are then plated in regeneration agar to

obtain transformants. The common methods for transforming *P. pastoris* are spheroplasting and PEG mediated transformations, lithium chloride mediated method, and electroporation. All method has its own advantages and disadvantages. These methods have two steps for the introduction of foreign DNA into *Pichia* cells; first is the preparation of competent yeast, and the second is the transformation of DNA into cells.

Previously we cloned and expressed *Xr* gene in pQE30Xa under the control of T7 promoter in *E. coli* M15 and used D-xylose and corncob during the fermentation for the production of xylitol in *E. coli* (Ariyan and Uthandi, 2019). *Pichia pastoris*, a methylotrophic yeast of ascomycetes class, is popularly used as a host to express heterologous proteins (Lin-Cereghino *et al.*, 2005). This is due to its growth to high cell densities, increased protein production (intracellular and extracellular), controlled and robust regulation of promoters, and easy scale-up (Ahmad *et al.*, 2014). Phillips Petroleum Company introduced *P. pastoris*, which utilizes methanol as a carbon source to produce single-cell protein as an additive to animal feed (Cregg *et al.*, 1985). The commonly used expression systems for protein production in *Pichia* is GAP (Glyceraldehyde-3-phosphate dehydrogenase) and AOX1 (alcohol oxidase) promoters. AOX1 promoter is widely used for controlled expression in which the gene of interest is placed between 5'AOX and 3'AOX (Vogl and Glieder., 2013). AOX1 and AOX2 genes

*Corresponding author's e-mail: usivakumartnau@gmail.com

synthesize alcohol oxidase, which utilize methanol and convert it to formaldehyde and hydrogen peroxide. AOX1 is responsible for 90 percent of activity.

AOX1 gets induced only when methanol is added; until then, it remains in a repressed state. Expression of AOX1 genes and protein production is restricted or lowered due to the presence of carbon sources other than methanol at the time of induction. Due to this, the cells are grown first in glycerol medium to attain growth and are transferred to methanol medium for induction and expression (Cereghino and Cregg, 2000). *P. pastoris* transformants selection system used in this study is based on auxotrophic selection. The wild strain is a deficit of *histidional dehydrogenase*, and so the vectors with yeast *his4* gene is used to transform the foreign DNA. As a result, the mutant *Pichia* can grow in a histidine deficient environment. Transformation efficiency is comparatively low in *P. pastoris* than in other yeast. Hence we performed three preferred methods of transformation for *P. pastoris* in order to select the best method to obtain transformants in *P. pastoris*.

In this paper, lithium chloride mediated, electroporation, and spheroplast methods were compared to transform *P. pastoris* GS115 with the *Xr* gene from *Candida tropicalis* cloned in vector pPIC3.5. The transformation protocol used, along with the results obtained for each method, is discussed.

MATERIAL AND METHODS

Strains, vectors, and media:

P. pastoris GS115 strain and vector pPIC3.5 were purchased from Invitrogen acquired by life technologies, USA (Catalog No. K1710-01). *P. pastoris* GS115 cells were grown in yeast extract peptone dextrose (YPD) broth (1% yeast extract, 2% peptone, 2% dextrose). The transformed cells were grown in regeneration dextrose medium (RDB) or minimal dextrose medium (MD). The vector pPIC3.5 containing *C. tropicalis XR gene* (*CtXr*) designated as pUSIXr was used (Fig 1). The plasmid was prepared using a plasmid DNA Kit (Thermo scientific Gene Jet plasmid miniprep kit, Ludhiana) followed by linearization with *SacI* and purified using spin columns (Qiagen, Germany). The purified linear DNA was quantified in nano-drop (Thermo Scientific, USA).

Spheroplasting method:

The procedure used to generate spheroplast were followed according to the protocol mentioned in the *Pichia* Expression Kit. A single pure colony of *P. pastoris* GS115 from a fresh plate was inoculated as a starter into 5 mL YPD broth and incubated at 30 °C overnight under shaking conditions at 120

rpm. The overnight grown culture was used for spheroplast preparation. About 5 µL of the overnight culture was used to inoculate 200 mL YPD broth grown in 500 mL conical flask. When the culture reached OD₆₀₀ of 0.2 to 0.3, the cells were harvested and washed in 20 mL of sterile water. The pellets were resuspended in SED solution (provided in the kit) (19 mL of SE + 1 mL of 1M DTT) and washed. About 20 mL of 1M sorbitol was used to wash cells. Finally, the cells were resuspended in 20 mL of SCE buffer (provided in the kit). About 10 mL of cells were used to determine the optimum time for spheroplasting by measuring OD at 800 nm in a multimode microplate reader (SpectraMax i3x, USA). Around 800 µL of 5% SDS and 200 µL of SCE were used as blank. The remaining cells were treated with zymolyase and incubated. The percentage of spheroplasting was calculated by using the formula

$$\% \text{ of spheroplasting} = 100 - (\text{OD}_{800} \text{ at time } t / \text{OD}_{800} \text{ at time } 0) \times 100$$

The second set of cells suspended in SCE buffer was added with 7.5 µL of zymolyase and incubated for a time at which 70% spheroplasting took place. After the optimum time, the cells were harvested by centrifugation at room temperature for 10 min at 750 x g. The spheroplast cells were washed with 10 mL of 1M sorbitol followed by 10 mL of CAS solution. Finally, the cells were suspended in 600 µL of CAS (provided in the kit). Spheroplast cells (100 µL) were taken in 2 mL of microfuge tubes, and 10 µg of linearized DNA was added to it and incubated at room temperature for 10 min, followed by the addition of 1 mL PEG/CaT solution and incubated further for 10 min at room temperature. Transformed cells were obtained by centrifugation at 750 x g for 10 min, and the pellets were resuspended in 150 µL SOS solution and incubated for 20 min. About 850 µL of 1 M sorbitol was added. From this, about 100 µL of cells were plated in RBD plates.

Lithium chloride method:

P. pastoris GS115 was grown in 50 mL of YPD to approximately an OD₆₀₀ of 1.0. The cells were harvested by centrifugation at 1500 x g and washed with 20 mL of sterile water. Cell pellets were resuspended in 1 mL of 100 mM lithium chloride and transferred to 1.5 mL microfuge tubes, and centrifuged at maximum speed for a few seconds. Supernatants were removed, and pellets were resuspended in 400 µL of 100 mM LiCl₂. About 50 µL of cells in lithium chloride suspension was used for single transformation. The cells were pelleted, and to that 240 µL of 50% PEG4000, 36 µL of 1M lithium chloride, 25 µL of carrier DNA (Salmon sperm DNA), and pUSIXr (5 µg) were added and vortexed approximately for 1 min. The tubes were incubated at 30 °C for 30 min, and heat shock was given at

42 °C for 25 min. Centrifugation was carried out at 8000 rpm, and the supernatant was removed. Pellets were resuspended in 1 mL of water, and from this, about 300 µL was used for plating. Plates were incubated at 30 °C for 2 days until the transformants appeared.

Electroporation method:

A single pure colony of *P. pastoris* GS115 was inoculated into 5 mL YPD broth and incubated overnight at 30 °C. Cells (1%) was inoculated into 200 mL of YPD. At an OD₆₀₀ of 1.3 to 1.5, the cells were harvested and resuspended in 500 mL of ice-cold sterile water. Cells were centrifuged and resuspended in 20 mL of ice-cold sorbitol. Cells were again centrifuged and finally suspended in 1 mL of 1M sorbitol. About 50 µL of competent cells were taken and mixed with 5 µg of linearized DNA. The mixture was transferred to 0.2 cm ice-cold electroporation cuvette and incubated in ice for 2 to 5 min. at 2000 kV of pulse was passed for 4 millisecc and immediately 1mL of 1M ice-cold sorbitol was added. About 300 to 500 µL of cells were used for plating and incubated for 4 to 6 days at 30 °C.

Screening of transformants:

The cells following transformation were plated in RDB or MD plates along with appropriate controls. The spheroplast cells were mixed with top agar containing RDB or MD followed by plating onto bottom RDB or MD plate. The viability of cells prepared during transformation was confirmed in RDBH or MDH. The transformed cells (~1 x 10²) were plated in RDB or MD plates after obtaining the transformants to confirm the growth, followed by colony PCR.

To confirm the presence of *Xr* gene in the transformants, selected colonies were subjected to colony PCR. Colonies from the plates were taken with a sterile toothpick, suspended in 10 µL of sterile water, and boiled at 95 °C for 10 min. The tubes were spun, and about 1 µL was used for 10 µL reaction. PCR reaction was performed using PCR Master Mix (Takara) with gene-specific *xr* primer and vector-specific *Aox* primers using the conditions of an initial denaturation at 94 °C for 5 min followed by 28 cycles of denaturation at 94 °C, annealing at 55 °C, and extension at 72 °C followed by a final extension at 72°C for 5 min. The PCR products were then run on 1% agarose gel to confirm the size of the product.

RESULTS AND DISCUSSION

Spheroplasting method:

In the present study, spheroplast was prepared by adding zymolyase to *P. pastoris* GS115. It was noticed that increasing the time of incubation after zymolyase addition increased the spheroplast

generation. The concentration of 70% spheroplast generated was used for transformation with 10 µg DNA. It was reported earlier that the transformation frequency increased when the concentration of spheroplasts was 3 x 10⁷– 3 x 10⁸. mL⁻¹, and the amount of single-stranded DNA were 1–3 µg. The frequency dramatically decreased when the concentration of spheroplasts was 2 x 10⁹.mL⁻¹ and the amount of DNA was 16 µg, indicating that the concentration of spheroplasts and amount of DNA were critical for high transformation frequency (Shigeyuki et al., 2010).




Figure 1. The map of cloned *C. tropicalis* *Xr* gene in pPIC3.5 plasmid (pUSIXr). The *Xr* gene was cloned in pPIC3.5 under the control of AOX promoter is indicated. The map was drawn using SnapGene software

After spheroplasting and transformation, the transformants were incubated at 28 °C and the transformants were observed on the second day of incubation. In our study, the spheroplast concentration of 70%, which had 3 x 10⁶ to 2 x 10⁷ was used with the DNA concentration of 10 µg. The colonies were subsequently selected on RBD plates to confirm the presence of the integrant gene. After selection, it was observed that less than 1 x 10⁴ to no colonies in the plate. Burgers and Percival (1987) proposed that decrease in efficiency of transformants may be due to high concentrations of lyticase to achieve rapid spheroplasting. Low transformation efficiency in the present study might be due to handling errors, loss of viability of cells during plating, and the proportion of DNA and cells during transformation (Burgers and Percival, 1987).

Lithium chloride method:

In intact *S. cerevisiae* cells, monovalent cation Li⁺ enhanced the transformation efficiency in the presence of PEG, and heat shock enhanced the transformation efficiency. Accordingly, we performed the transformation of *P. pastoris* GS115 with 5 µg of pUSIXr. The transformants were selected on MD plates. The transformants appeared on the 2nd or

3rd day of incubation with an average transformation efficiency of 2×10^3 . Ito et al. (1983) obtained the transformation efficiency of 4×10^2 using $1 \mu\text{g}$ of DNA by treating the cells with lithium chloride. The procedure included plasmid DNA and polyethylene glycol (PEG), followed by a heat shock at 42°C .

Around 30 to 40 % of competent cells take up one or more plasmid molecules, so the increase in plasmid concentration increases efficiency. Tsai et al., (2017) obtained fewer transformants ($20 \mu\text{g}$ of DNA) by following lithium acetate PEG mediated transformation methods. Comparatively, in the




Figure 2. Generation of spheroplast using enzyme zymolyase: Spheroplasting was generated by incubating the cells after the addition of zymolyase and extending the time of incubation until the desired spheroplast was obtained. The OD800 (primary axis) denotes the measure of viable cells and the spheroplasting (%) (secondary axis) increases with time is indicated. The optimal spheroplasting (70%) was obtained during 25 minutes of incubation with zymolyase.

present study, the lithium chloride method has given efficient transformation per μg pUSIXr.

Electrophoration method:

The Electrophoration method is a widely used efficient method for transformation. In our study, we used electrotransformation employing 1500 KV to 2000 KV for 5 ms with $5 \mu\text{g}$ pUSIXr. After giving the pulse, the cells were immediately chilled by the addition of 1M sorbitol followed by plating. The transformants were obtained on the 4th day of plating in RDB or MD. The transformation efficiency obtained was higher when compared with other methods of transformation. An average of 4×10^5 to 1×10^6 cells were obtained using this method (Wu and Letchworth, 2004) obtained 30–50 colony forming units (CFU) per microgram DNA at 2.5

to 3.5kv for 4–5 ms. The highest transformation efficiency in *P. pastoris* was obtained at 1.5 kV, $25 \mu\text{F}$, and 186Ω with 2.6×10^5 to 4×10^6 per μg DNA with pPIC9K. A maximum of 2.6×10^5 transformants was produced when $1 \mu\text{g}$ of pPIC9K DNA was used (Wu and Letchworth, 2004).

Transformation efficiency in *P.pastoris* GS115:

Transformation of *P. pastoris* GS115 using spheroplast, LiCl₂, and electroporation resulted in more recombinants in LiCl₂ method followed by electroporation. The colonies appear in the selection plate after 2 to 3 days of incubation in lithium chloride method, and it took 4 to 6 days for electroporation and spheroplast methods. The recombinants were confirmed by plating in RBD plate and control (Fig 3) and also by using a gene-specific




Figure 3: Representative transformants grown on YPD (1) and RDB (2) plate. *P. pastoris* GS115 wild type and transformants were streaked on YPD and RDB plates. The recombinants were able to synthesis His and were able to grow on RDB plate, whereas wild type *P. pastoris* GS115 did not grow on RDB plate. Both recombinants and wild type could grow on PDB control plate. The recombinants were positive for Xr gene amplification.

primer PCR to confirm the integrated gene. Among the three methods, maximum transformants were obtained in the electroporation followed by lithium chloride for GS115, proving electroporation and LiCl₂ as a method of choice suitable for transforming *P. pastoris* GS115.

Acknowledgment

Authors acknowledge the financial assistance from University Core Project (TNAU-Core Project-CP016) to SU and Indo- Russia Joint collaboration supported by DBT, Gol, New Delhi (No.DBT/IC2/ Indo-Russia/2014-16/04) to SU. We are also thankful to DBT-BIOCARe (No. BT/PR18134/BIC/101/795/2016) for the grant awarded to PR. The authors are grateful to the instrumentation facility extended by the metabolomics and proteomics analytical facility, Dept. of Agrl. Microbiology, TNAU, established under Gol-MHRD (FAST)-scheme.

Author contributions:

Research grant and Idea conceptualization, US; Experiments, PY; Guidance, US, RP Writing-original draft, PY; Writing- reviewing & editing, US, RP

REFERENCES

Ahmad, M., M. Hirz, H. Pichler, and H. Schwab. 2014. "Protein expression in *Pichia pastoris*: recent achievements and perspectives for heterologous protein production." *Applied Microbiology and Biotechnology* 98(12): 5301-5317.

Ariyan, M., and Uthandi, S. 2019. "Xylitol Production by Xylose Reductase over producing Recombinant *Escherichia coli* M15." *Madras Agric. J.* 106 (1-3): 1.

Burgers, P.M.J., and K.J. Percival. 1987. "Transformation of yeast spheroplasts without cell fusion." *Analytical Biochemistry* 163(2): 391-397.

Cereghino, J.L., and J.M. Cregg. 2000. "Heterologous protein expression in the methylotrophic yeast *Pichia pastoris*." *FEMS Microbiology reviews* 24(1): 45-66.

Cregg, J.M., K.J. Barringer, A.Y. Hessler, and K.R. Madden. 1985. "*Pichia pastoris* as a host system for transformations." *Molecular and Cellular Biology* 5(12): 3376-3385.

Hinnen, A., J.B. Hicks, and G.R. Fink. 1978. "Transformation of yeast." *Proceedings of the National Academy of Sciences* 75(4): 1929-1933.

Ito, H., Y. Fukuda, K. Murata, and A. Kimura. 1983. "Transformation of intact yeast cells treated with alkali cations." *Journal of Bacteriology* 153(1): 163-168.

Jung, I.-Y., J.-W. Lee, W.-K. Min, Y.-C. Park, and J.-H. Seo. 2015. "Simultaneous conversion of glucose and xylose to 3-hydroxypropionic acid in engineered *Escherichia coli* by modulation of sugar transport and glycerol synthesis." *Bioresource technology* 198: 709-716.

Kawai, S., W. Hashimoto, K. Murata. 2010. "Transformation of *Saccharomyces cerevisiae* and other fungi: methods and possible underlying mechanism." *Bioeng Bugs* 1(6): 395-403.

Lin-Cereghino, J., W.W. Wong, S. Xiong, W. Giang, L.T. Luong, J. Vu, S.D. Johnson, and G.P. Lin-Cereghino. 2005. "Condensed protocol for competent cell preparation and transformation of the methylotrophic yeast *Pichia pastoris*." *Biotechniques* 38(1): 44-48.

Mohamad, N.L., S.M. Mustapa Kamal, and M.N. Mokhtar. 2015. "Xylitol biological production: a review of recent studies." *Food reviews international* 31(1): 74-89.

Orr-Weaver, T.L., J.W. Szostak, and R.J. Rothstein. 1981. "Yeast transformation: a model system for the study of recombination." *Proceedings of the national Academy of Sciences* 78(10): 6354-6358.

Tsai, Y.-Y., T. Ohashi, T. Kanazawa, P. Polburee, R. Misaki, S. Limtong, and K. Fujiyama. 2017. "Development of a sufficient and effective procedure for transformation of an oleaginous yeast, *Rhodospiridium toruloides* DMKU3-TK16." *Current genetics* 63(2): 359-371.

Wu, S., and G.J. Letchworth. 2004. "High efficiency transformation by electroporation of *Pichia pastoris* pretreated with lithium acetate and dithiothreitol." *Biotechniques* 36(1): 152-154.