



RESEARCH ARTICLE

Optimization of Copper Sulphate Levels to Enhance Yield and Quality of Aggregatum Onion (*Allium cepa* var *aggregatum* L)

D. Jegadeeswari*, T. Chitdeshwari* and A.K. Shukla** 1 *

Department of Soil Science and Agricultural chemistry Tamil Nadu Agricultural University Coimbatore - 641 003
Indian Institute of Soil Science, Bhopal -462 038.

ABSTRACT

Field experiments were conducted during rabi season of 2018 and summer season of 2019 at farmer's field, western zone of Tamil Nadu at Pattiyarkovilpathy, Theethipalayam and Narasipuram villages of Thondamuthur block of Coimbatore district. Totally five treatments comprising of varied CuSO_4 levels (0, 2.5, 5.0, 7.5 & 10.0 kg ha^{-1}) were replicated four times in a randomized block design under irrigated conditions. The results showed that, fresh mean bulb yield of onion in four locations ranged from 12.4 to 15.6 t ha^{-1} . The highest mean bulb yield of 15.6 t ha^{-1} was registered with recommended NPK+5 $\text{kg CuSO}_4 \text{ ha}^{-1}$ and the lowest mean bulb yield of 12.4 t ha^{-1} was recorded in NPK control with no CuSO_4 addition. Dry Matter Production ranged from 1065 to 1199 kg ha^{-1} . The highest DMP of 1199 kg ha^{-1} was recorded in the treatment that was applied with NPK+5 $\text{kg CuSO}_4 \text{ ha}^{-1}$ and the lowest DMP (1065 kg ha^{-1}) was noticed with NPK control. Concerning copper content in bulb and foliage, it ranged from 6.09 to 12.3 and 8.46 to 13.7 mg kg^{-1} with the location mean of 6.16 to 12.0 mg kg^{-1} and 7.61 to 11.1 mg kg^{-1} , respectively. Copper uptake in bulb and sheath varied from 7.05 to 18.3 and 6.26 to 11.0 g ha^{-1} (mean of 8.55 to 12.3 and 6.52 to 10.4 g ha^{-1}), respectively with a total uptake of 14.1 to 27.6 g ha^{-1} . In both bulb and foliage, Cu content and uptake, increased with increasing doses of CuSO_4 and the highest Cu content and uptake was registered with the application of CuSO_4 @ 10.0 kg ha^{-1} . Regarding the availability of copper in soil, it ranged from 0.73 to 2.42 mg kg^{-1} and increasing levels of CuSO_4 application increased the DTPA Cu status in soil and application of 10.0 $\text{kg CuSO}_4 \text{ ha}^{-1}$ recorded the highest copper availability in soil. Quality parameters viz., TSS, titrable acidity and ascorbic acid varied from 10.25 to 14.08 °Brix, 0.28 to 0.61% and 12.9 to 20.1 (100 g^{-1}FW), respectively. Field experiments conducted to optimize the rates of copper sulfate application to improve the yield and quality of aggregatum onion in four locations at Coimbatore district, showed that, soil application of 5 kg CuSO_4 along with recommended NPK @ 60:30:30 kg ha^{-1} was found optimal for obtaining higher bulb yield (15.6 t ha^{-1}) and better quality in aggregatum onion with a BC ratio of 4.63.

Keywords: Aggregatum onion - copper sulphate - optimization- -yield, quality

INTRODUCTION

Onion (*Allium cepa* L.), belonging to the family Alliaceae, is one of the most important and popular vegetable spice crops cultivated worldwide (Mishra *et al.*, 2013). It is famous for its characteristic flavour and is widely used to increase the taste of foods like gravies, soups, stew stuffing, fried fish, and meat (Rashid *et al.*, 2016). The increased productivity with high quality onion is the utmost demand of the onion growers. Lack of micronutrients addition in the balanced fertilization schedule

is one of the important causes for low yield and quality (Lata Shukla *et al.*, 2015). Onion requires a sufficient amount of plant nutrients and responds very well to the added nutrients (Alam *et al.*, 2010). Therefore, systematic fertilization of different micronutrients for onion cultivation is needed for the low organic matter and micronutrient deficient soils (Lata Shukla *et al.*, 2015). Onion is more susceptible to nutrient deficiencies than most crop plants because of their shallow and unbranched root system; hence they require and often respond

*Corresponding author's e-mail: elaento@gmail.com

well to the addition of fertilizers (Brewster, 1994). Among the various factors affecting the yield of onion, adequate and balanced nutrient management plays a major role to optimise the quality and quantity of harvested plant products (Lakshmi and Sekhar, 2018). Nutrient management involves using crop nutrients as efficiently as possible to improve productivity while protecting the environment. The key fundamental principle behind nutrient management is balancing soil nutrient inputs with crop requirements (Khalid Mahmud Khokhar, 2019). Among all the essential nutrients, micronutrients viz., Fe, Mn, Zn, Cu, B and Mo are important for the optimal growth and productivity of the plants. Though these elements are required in small amounts, they are very important for plant development and for profitable crop production because they act as activators of many plant functions. Micronutrients among essential elements, plays a catalytic role in nutrient absorption and balancing other nutrients (Singh and Kalloo, 2000, Poongothai and Chitdeshwari, 2003). Although the requirement of a micronutrient is small compared to macronutrients, nevertheless, micronutrient deficiency can limit crop growth and production. Furthermore, micronutrients help increase the efficiency of the use of macronutrients. Micronutrients have a great role in the fertilizer program to achieve higher and sustainable crop yields (Vijay Aske et al., 2017 and Jegadeeswari et al., 2019). Among the micronutrients, copper is important element for plant growth because it acts as a catalyzer of several plant processes, imparts disease resistance in plants, indirect role in chlorophyll production, intensifies the color and flavor of fruits and vegetables and playing an irreplaceable role in the function of a large number of enzymes which catalyze oxidative reactions in a variety of metabolic pathways. Cu can also be considered as a toxic element whose deleterious effects usually arise at high concentrations (Donghua Liu et al., 1994). The copper deficiency interferes with protein synthesis and causes a buildup of soluble nitrogen compounds. Normal plants contain 8 to 20 mg kg⁻¹ copper and deficient plants usually contain less than 6 mg kg⁻¹. Copper deficiency in many plants shows

up as wilting or lack of turgor and development of a bluish green tint before leaf tips become chlorotic and die. Onion bulbs show poor pigmentation because of copper deficiency (El-Hadidi et al., 2016). Looking at these facts, the present research was undertaken to optimize the dose of copper sulfate for increasing the productivity and quality of onion.

MATERIAL AND METHODS

Field experiment and laboratory analysis

Four field experiments were conducted, two each during rabi season of 2018 and two during summer season of 2019 at farmer's fields, western zone of Tamil Nadu at Pattiyarkovilpathy, Theethipalayam and Narasipuram villages of Thondamuthur block, Coimbatore district, Tamil Nadu, India. Initial soil sample was collected from 0-15 cm depth and analyzed for various physico-chemical and chemical characteristics (Table 1). Onion variety local was used as a test crop and cropping system was irrigated.

Table 1. Details of analytical procedures employed in soil and plant sample analysis

Parameters	Methodology	References
Physico-chemical Properties Soil reaction (pH)	Potentiometry (1:2.5 soil : water suspension)	(Jackson, 2005)
Electrical conductivity	Conductometry (1:2.5 soil: water suspension)	(Jackson, 2005)
Chemical Properties Soil Organic Carbon Available nutrients	Chromic acid wet digestion method	(Wakley and Black. 1934)
Nitrogen (N)	Alkaline permanganate method	(Subbiah and Asija, 1956)
Phosphorus (P)	Olsen's method	(Olsen et al., 1954)
Potassium (K)	Neutral N NH ₄ OAc	(Stanford and English, 1949)
DTPA Zn, Cu, Fe and Mn Plant sample analysis	DTPA extraction, AAS method	(Lindsay and Norvell, 1978)
Cu	Triple acid digestion and AAS method	(Lindsay and Norvell, 1978)
Quality parameters	-	Ranganna, S. 1997

Experimental design

Totally five treatments comprising of varied Cu levels (0, 2.5, 5.0, 7.5 & 10.0 kg CuSO₄ ha⁻¹) were replicated four times in a randomized block design. The recommended dose of NPK fertilizers at

30:60:30 was applied as basal, 30 kg N alone was applied as a top dressing on 30 DAS to all the treatments. Urea as N source, Single super phosphate for P, Muriate of Potash for K and Copper sulfate for Cu were used as inorganic fertilizer sources.

Table 2. Initial Soil characteristics

L. No	pH	EC (dS m ⁻¹)	OC (%)	Major nutrients (kg ha ⁻¹)			S (mg kg ⁻¹)	Micronutrients (mg kg ⁻¹)				
				N	P	K		Fe	Mn	Cu	Zn	B
1	7.3	0.40	0.45	204	15.8	288	22.3	15.5	7.8	0.32	0.60	0.56
2	7.40	0.31	0.40	185	16.8	309	26.0	12.5	6.3	0.66	0.74	0.88
3	7.0	0.42	0.54	154	14.4	240	28.2	13.1	6.8	0.57	0.81	0.74
4	7.7	0.50	0.60	160	18.1	275	20.3	16.3	8.4	0.31	0.63	0.71

The crop was grown to maturity with routine cultural operations by imposing the treatments and at harvest bulb yield, dry foliage yield, yield attributing characters and quality parameters like total soluble sugars, ascorbic acid and titrable acidity were recorded besides collecting plant and post-harvest soil samples for analyzing the Cu availability, its content and uptake in bulb and leaves.

Statistical analysis

The experimental data collected relating to different parameters were statistically analyzed using the procedure given by Panse and Sukhatme, 1967. The data were subjected to Fisher's method of analysis of variance and the level of significance used in F test was $P = 0.05$. The critical difference was calculated at 5 per cent probability level whenever F value was found to be significant.

RESULTS AND DISCUSSION

The soil reaction of the experimental initial soil was neutral in all the locations with pH values of 7.30, 7.40, 7.0 and 7.7 with nil salinity status having

gets decreased. Highest bulb yield of 14.6, 15.3, 16.4 and 16.0 t ha⁻¹ was registered in Location 1, 2,3 & 4, respectively with the a mean of 15.6 t ha⁻¹. It was followed by 7.5 kg CuSO₄ ha⁻¹ in all the locations with the mean yield of 15.0 t ha⁻¹ and the lowest bulb yield of 11.8, 12.6, 13.5,11.5 t ha⁻¹ with a mean of 12.4 t ha⁻¹ was recorded in the NPK control with no CuSO₄ in four locations (Table.3 & Fig.1).


Table 3. Effect of different levels of CuSO₄ on the yield, dry matter production and BCR

Levels of CuSO ₄ (kg ha ⁻¹)	Bulb yield (t ha ⁻¹)					DMP (kg ha ⁻¹)					BCR
	L1	L2	L3	L4	Mean	L1	L2	L3	L4	Mean	
0	11.8	12.6	13.5	11.5	12.4	1093	1031	1100	1036	1065	3.72
2.5	13.6	13.9	14.2	12.9	13.7	1123	1059	1127	1063	1093	4.09
5.0	14.6	15.3	16.4	16.0	15.6	1227	1163	1241	1165	1199	4.63
7.5	14.3	14.7	15.8	15.2	15.0	1163	1099	1180	1105	1137	4.43
10.0	14.0	14.2	15.4	13.9	14.4	1109	1049	1132	1057	1087	4.24
Mean	13.7	14.1	15.1	13.9		1143	1080	1156	1085		
SEd	0.26	0.17	0.16	0.30		32.6	33.6	30.2	30.9		
CD (P= 0.05)	0.58	0.37	0.35	0.67		76.1	78.0	69.4	71.7		

Similar A similar trend was observed with respect to the dry matter production of onion in four locations and the DMP in all the locations ranged from 1031 to 1241 kg ha⁻¹ with a mean range of 1065 to 1199

kg ha⁻¹. The highest DMP of four locations were 1227,1163,1241,1165 kg ha⁻¹ with a mean of 1199 kg ha⁻¹, which was recorded in the treatment that was applied with NPK+5 kg CuSO₄ ha⁻¹ and the lowest

Figure.1. Optimizing the rate of copper dose for onion



the electrical conductivity of 0.40, 0.31, 0.42 and 0.50 dSm⁻¹. The organic carbon content was 0.45, 0.40, 0.54 and 0.60 g kg⁻¹ in all four locations. Available nitrogen and phosphorus contents of the soils were low 204, 185, 154, 160 kg ha⁻¹ and 15.8, 16.8, 14.4 and 18.1 kg ha⁻¹ respectively. The values for available K and S were 288, 309, 240, 275 kg ha⁻¹ and 22.3, 260, 28.2 and 20.3 mg kg⁻¹ respectively. The nutrient status of all the farm soils showed low to medium in organic carbon content, low to medium in available N and P, high in available K and S. Micronutrient availability in the soils revealed sufficient status of Mn and Fe, while many of the farms were deficient in Zn and Cu availability. DTPA Copper status in the soils were 0.72, 0.66, 0.57 and 0.81 mg kg⁻¹. The experimental soil was sandy loam.

Yield and DMP

The results showed that, fresh bulb yield of onion in all four locations ranged from 11.5 to 16.4 t ha⁻¹ and a mean variation of 12.4 to 15.6 t ha⁻¹. Application of CuSO₄ at graded levels increased the onion bulb yield up to 5 kg CuSO₄ ha⁻¹ and thereafter

DMP of 1093,1031,1100,1036 kg ha⁻¹ with a mean of 1065 kg ha⁻¹ was noticed with NPK control with no CuSO₄ (Table.3). Response to CuSO₄ was found to be very high in the soils having deficient Cu availability. This could be attributed to the vital functions of copper in plant growth and metabolism. Application of CuSO₄ at higher levels up to 5 kg per ha resulted in the development of good growth of sheath, bulbs and accelerated protein synthesis, bulb formation and better yield of crop. The results are further in accordance with the reports of Chattopadhyay et al. (2004) and Ballabh et al., (2013), who reported that the yield characters are significantly influenced with a higher level of micronutrients like, Zn, B, Mn

and Cu in onion. Bose *et al.*, (2009) stated that the increase in growth and yield might be due to role of copper as essential for plant metabolism, one of the constituents of certain oxidizing reduction enzymes. Therefore, its role in plant metabolism and participation in oxidation-reduction action, check in certain diseases and there by improved growth of plant and yield. The benefit benefit-cost ratio ranged from 3.72 to 4.63, a maximum of 4.63 was observed in the treatment with the application of NPK + 5.0 kg CuSO₄ ha⁻¹ and the lowest was noticed in NPK control with no CuSO₄ (Table.3).

Cu content and uptake Copper

The copper content in both onion bulb and foliage was estimated and the uptake was computed. The results showed that application of copper sulfate resulted in increased Cu content in onion bulb and sheath and it was increased with increasing levels of CuSO₄. In onion bulb, copper content in all four locations varied from 6.09 to 12.3 mg kg⁻¹ with a mean variation of 6.16 to 12.0 mg kg⁻¹. In all the locations, the mean value showed the highest Cu content of 12.3,11.4,11.9,12.2 mg kg⁻¹ with an

overall mean of 12.0 mg kg⁻¹ in the treatment that received NPK+10 kg CuSO₄ ha⁻¹ which was followed by 7.5 kg CuSO₄ ha⁻¹ addition and the lowest mean values was noticed in the NPK control. With regard to copper content in foliage, the values ranged from 8.46 to 13.7 mg kg⁻¹ with the location mean of 7.61 to 11.1 mg kg⁻¹ (Table 4). A similar trend was observed in the foliage also with the increased values for the increased levels of

Table 4. Effect of different levels of CuSO₄ on the Cu content in onion

Levels of CuSO ₄ (kg ha ⁻¹)	Cu content (mg kg ⁻¹)									
	Bulb					Foliage				
	L1	L2	L3	L4	Mean	L1	L2	L3	L4	Mean
0	6.09	6.17	6.24	6.13	6.16	8.65	8.57	8.51	8.46	8.55
2.5	7.19	6.94	7.24	6.82	7.05	9.17	9.21	9.09	9.27	9.19
5.0	8.73	8.59	8.67	8.49	8.62	10.9	11.1	11.0	11.3	11.1
7.5	9.95	9.79	9.88	9.69	9.83	11.6	12.4	11.8	12.7	12.1
10.0	12.3	11.4	11.9	12.2	12.0	13.1	13.4	12.9	13.7	13.3
Mean	8.85	8.58	8.79	8.67	8.67	10.7	10.9	10.7	11.1	10.8
SEd	0.26	0.17	0.16	0.30	0.11	0.11	0.21	0.16	0.07	0.15
CD (P= 0.05)	0.58	0.37	0.35	0.67	0.24	0.46	0.35	0.15	0.15	0.25

CuSO₄ application. In all the locations, the highest copper content of 13.1,13.4,12.9,13.7 mg kg⁻¹ and a mean 13.3 mg kg⁻¹ was recorded in the treatment applied with NPK + CuSO₄@ 10.0 kg ha⁻¹ which was

followed by 7.5 kg CuSO₄ addition and the values were 11.6,12.4,11.8,12.7 mg kg⁻¹ with a mean of 12.1 mg kg⁻¹. Similarly, the lowest copper content of 8.65, 8.57, 8.51, 8.46 mg kg⁻¹ and a mean of 8.55 mg kg⁻¹ were observed with NPK control with

no CuSO₄. Copper uptake in bulb and sheath also increased with increasing dose of copper sulphate and the values varied from 7.05 to 18.3 and 6.26 to 11.0 g ha⁻¹ (mean of 8.55 to 12.3 and 6.52 to 10.4 g ha⁻¹) respectively. In both bulb and foliage,

the highest uptake was registered with the application of CuSO₄ @ 10.0 kg ha⁻¹. In all the four locations, in bulb the highest Cu uptake of 17.2,16.2,18.3,17.0 g ha⁻¹ with a mean of 17.2 g ha⁻¹ (Table 5) was recorded with NPK+10 kg ha⁻¹ CuSO₄ addition, which was followed by NPK+7.5 kg ha⁻¹ CuSO₄ addition and the values were


Table 5. Effect of different levels of CuSO₄ on the Cu uptake by onion

Levels of CuSO ₄ (kg ha ⁻¹)	Cu content (mg kg ⁻¹)									
	Bulb					Foliage				
	L1	L2	L3	L4	Mean	L1	L2	L3	L4	Mean
0	7.19	7.77	8.42	7.05	7.61	6.76	6.78	6.26	6.29	6.52
2.5	9.78	9.65	10.28	8.80	9.63	7.44	7.53	6.94	7.15	7.27
5.0	12.7	13.1	14.2	13.6	13.4	9.9	10.4	9.5	9.9	9.94
7.5	14.2	14.4	15.6	14.7	14.7	10.0	10.8	9.5	10.3	10.1
10.0	17.2	16.2	18.3	17.0	17.2	10.4	11.0	9.7	10.5	10.4
Mean	12.2	12.2	13.4	12.2	12.2	8.90	9.30	8.38	8.83	8.83
SEd	0.29	0.37	0.28	0.34	0.14	0.14	0.21	0.21	0.21	0.21
CD (P= 0.05)	0.63	0.82	0.61	0.75	0.32	0.47	0.46	0.27	0.27	0.32

14.2,14.4,15.6,14.7 g ha⁻¹ with a mean of 14.7 g ha⁻¹. In bulb, the lowest copper uptake of 7.19,7.77,8.42,7.05 g ha⁻¹ with a mean of 7.61 g ha⁻¹ which was recorded in the NPK control with no CuSO₄. Similarly, in foliage, also

similar trend was observed and in all four locations, the highest Cu uptake of 10.4,11.0,9.7,10.5 g ha⁻¹ with a mean of 10.4 g ha⁻¹ was recorded with NPK+10 kg ha⁻¹ CuSO₄ addition, which was followed by NPK+7.5 kg ha⁻¹ CuSO₄ and the values were 10.0,10.8,9.5,10.3 g ha⁻¹ with a mean of 10.1 g ha⁻¹. In foliage, the lowest copper uptake of 6.76,6.78,6.26,6.29 g ha⁻¹ with a mean of 6.52 g ha⁻¹ was recorded in the NPK control.

Figure.2. Effect of Copper sulphate application on DTPA Cu availability in soil



However, the uptake of plant nutrients depends on several factors, such as cultivar, crop environment, soil fertility, fertilization methods (Lee *et al.*, 2009; Yoldas *et al.*, 2011).

Quality parameters

The quality parameters of onion were significantly influenced by the application of copper sulfate at

1.50, 1.24, 1.30, 5.65 42 mg kg⁻¹ with a mean of 2.42 42 mg kg⁻¹ in soil. The copper availability in post harvest soil showed the sustained status of copper except NPK control, which recorded the lowest value of 0.73 mg kg⁻¹.

Table 6. Effect of different levels of CuSO₄ on quality parameters of onion

Levels of CuSO ₄ (kg ha ⁻¹)	TSS					Ascorbic acid					Titrate acidity				
	L1	L2	L3	L4	Mean	L1	L2	L3	L4	Mean	L1	L2	L3	L4	Mean
0	11.0	10.3	11.6	11.1	10.9	13.1	14.6	12.9	16.5	14.3	0.32	0.48	0.37	0.43	0.40
2.5	12.1	12.9	12.6	12.9	12.7	15.2	16.9	14.1	17.9	16.1	0.34	0.56	0.38	0.50	0.45
5.0	12.4	13.4	14.1	13.9	13.5	16.5	18.9	17.9	20.1	18.4	0.36	0.61	0.48	0.52	0.49
7.5	12.1	11.9	13.1	12.3	12.4	16.4	17.1	16.6	17.4	16.9	0.34	0.44	0.34	0.48	0.40
10.0	11.1	11.7	11.9	10.8	11.4	13.7	14.1	14.5	16.0	14.6	0.30	0.42	0.28	0.35	0.34
SEd	0.96	1.00	0.27	0.19	0.51	0.51	0.42	0.55		0.06	0.05	0.07	0.02		
CD (P= 0.05)	NS	NS	0.60	0.43		1.12	1.13	0.92	1.22		NS	0.10	NS	0.04	

This might be due to the effect of the addition of Cu through fertilisers and also available from inherent properties of soil like, parent material, the effect of organic matter and presence of clay minerals. Similar findings was reported by Kamble and Kathmale (2015).

CONCLUSION

Relevant Field experiments were conducted to optimize the dose of copper sulfate for aggragaum onion to increase the yield and quality under irrigated conditions. The highest bulb yield of 15.6 t ha⁻¹ with the addition of recommended NPK+5 kg CuSO₄ ha⁻¹ and the lowest bulb yield of 12.4 t ha⁻¹ was recorded in NPK control. Mean DMP ranged from 1065 to 1199 g ha⁻¹. In addition, the application of copper

sulfate increased the Cu content both in sheath and bulb thus increased Cu uptake was noticed with increasing levels of CuSO₄ and the highest content and uptake was recorded with NPK+10 kg CuSO₄ ha⁻¹. Regarding the availability of copper in the soil, the availability was found to be sufficient at all levels of CuSO₄ addition. Increasing levels of copper application increased the soil Cu status and application of 10.0 kg CuSO₄ ha⁻¹ recorded the highest copper availability in soil. Application

Cu availability

Regarding the availability of copper in soil, it ranged from 0.73 to 2.42 mg kg⁻¹ and the availability was found to be sufficient at all levels of CuSO₄ addition (Fig.2). Increasing levels of application increased the soil Cu status and in all four locations, application of 10.0 kg CuSO₄ ha⁻¹ recorded the highest DTPA Cu of

of CuSO_4 @ 5 kg ha⁻¹ also registered better quality attributes viz., total soluble solids, ascorbic acid and titrable acidity in onion bulb. The above field experiments conducted to optimize the rates of copper sulfate application to improve the yield and quality of aggregatum onion in four locations at Coimbatore district, showed that, soil application of 5 kg CuSO_4 along with recommended NPK @ 60:30:30 kg ha⁻¹ was found optimal for obtaining higher bulb yield (15.6 t ha⁻¹) and better quality in aggregatum onion with a BC ratio of 4.63.

ACKNOWLEDGMENT

The authors greatly acknowledge the financial support rendered by the AICRP-Micronutrients Scheme of Indian Society of Soil Science, Bhopal and Tamil Nadu Agricultural University, Coimbatore for the execution of technical programme, man power and facility support.

REFERENCES

- Acharya, N., Venkatesan, K., Saraswathi, T and K.S. Subramanian. 2015. Effect of Zinc and boron application on growth and yield parameters of multiplier onion (*Allium cepa* L. var. aggregatum Don) var. Co 5. *International Journal of Research*. 2(1): 757-765
- Alam, M.N., Abedin, M.J. and M.A.K. Azad. 2010. Effect of micronutrients on growth and yield of onion under calcareous soil environment. *International Research Journal of Plant Science* 1(3): 056-061.
- Aske V, Jain PK and S. Govind. 2017. Effect of Micronutrients on yield, quality and storability of onion cv. Bhima Super. *Trends in Biosciences*. (10):1354-1358. 9
- Ballabh, K., Rana, D.K. and S.S. Rawat. 2013. Effects on foliar application of micronutrients on growth, yield and quality of onion. *Indian Journal of Horticulture*, 70(2): 260-265.
- Brewster, J.L. 1994. Onions and other vegetable Alliums. CAB International, Wallingford, UK, p. 36.
- Chattopadhyay, S.B. and T.P. Mukhopadhyay. 2004. Response of boron and molybdenum as foliar feeding on onion in tarai soil of West Bengal. *Environment and Ecology*, 22(4): 784-787.
- Donghua Liu, Wusheng Jiang, Cheng Lu, Fengmei Zhao, Yuqing Ha0 and Lin Guo. 1994 Effects of copper sulfate on the nucleolus of *Allium cepa* root-tip cells. *Heredilus*, 120: 87-90
- El-Hadidi, E.M. ; M.M. El-Shazly and Heba M.M. Hegazy. 2016 Effect of N, P and Cu fertilization on onion yield, quality and nutrients uptake . *J. Soil Sci. and Agric. Eng., Mansoura Univ.*, Vol. 7(2): 231 - 236
- Fink M., Feller C., Scharpf H.C., Weier U., Maync A., Ziegler J., Paschold P.J., and K. Strohmeyer. 1999. Nitrogen, phosphorus, potassium and magnesium contents of field vegetables – recent data for fertilizer recommendations and nutrient balances. *Journal of Plant Nutrition and Soil Science*, 162: 200 71-73.
- Jackson, M. L. 2005. *Soil chemical analysis: Advanced course (2-Ed)* UW-Madison Libraries, Madison, Wisconsin, p. 798-837. <http://parallelpress.library.wisc.edu>
- Jegadeeswari, D, Chitdeshwari, T and P. Boominathan. 2019. Screening of short duration rice genotypes for Zn use efficiencies. *International J. of Experimental Biology and Agricultural Sciences*, 7(1): 25- 205 33
- Kamble, B.M and D.K. Kathmale. 2015. Effect of different levels of customized fertilizer on soil nutrient availability, yield and economics of onion. *J. of Applied and Natural Sci.*, 7(2):817-821
- Lakshmi, C.S and C.C. Sekhar. 2018. A review on integrated nutrient management in bulbous vegetable crops. *Journal of Pharmacognosy and Phytochemistry*, 7, 411-415
- Lata Shukla, U.S. Bose and Manoj Kumar Ahirwar. 2015. Effect of foliar feeding of micronutrients on growth, yield and income from rabi onion var. Agrifound light red. *Annals of Plant and Soil Research* 17 (3): 307-310
- Lee J., Lee S. 2014. Correlations between soil physicochemical properties and plant nutrient concentrations in bulb onion grown in paddy soil. *Scientia Horticulturae*, 179: 158-162.
- Lee E.J., Yoo K.S., Jifon J and B.S. Patil. 2009. Application of extra sulphur to high-sulphur soils does not increase pungency and related compounds in shortday onions. *Scientia Horticulturae*, 123: 178- 217 183.
- Lindsay, W. L. and W.A. Norvell. 1978. Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil science society of America journal*, 42 (3), 421-428.
- Mishra, H.P., Sarkar, C., Viswajith, K.P., Dhekale, B.S. and P.K. Sahu. 2013. Instability and forecasting using ARIMA model in aua, Production and productivity of onion in India. *Journal of Crop and Weed*, 9: 96- 222 101
- Olsen, S. R., Cole, C. V., Watanabe, F. S. and L.A. Dean. 1954. Estimation of available phosphorus in soil by extraction with sodium bicarbonate. *U. S. Dept. Agri, Circ.*, 939.
- Panse, V.G. and P.V. Sukhatme. 1967. *Statistical methods for agricultural workers*, ICAR Publications, New Delhi.
- Poongothai. S and T. Chitdeshwari. 2003. Response of blackgram to multi-micronutrients, *Madras Agricultural Journal*, 90, 442-443
- Rashid, M.H.A, Massiah, A.J. and Thomas, B. 2016. Genetic regulation of day length adaptation and bulb formation in onion (*Allium cepa* L.). *Acta Horticulturae*, 1143: 7-14
- Ranganna, S. 1997. *Hand Book of analysis and quality control for fruits and vegetables products*. 3, & edition, Tata McGraw - Hill publishing company Ltd. New Delhi, 12-16
- Rupali Dogra, Ashok K. Thakur and Uday Sharma. 2019. Studies on Agronomic Bio-fortification with Micronutrients in Onion (*Allium cepa* L.) Seed Crop. *International Journal of Economic Plants*. 235 6(3):136-139

- Singh, K.P and G. Kalloo. 2000. Nutrient management in vegetable crops. Fertilizer News 45:77–81.**
- Subbiah, B and G.L.Asija. 1956.Alkaline Permanente method of available nitrogen determination. Current Science, 25 : 259-260**
- Tariq A Bhat, Chattoo, MA, Parray, EA, Mushtaq,F, Asif M Rather, Zehra, SB, Hajam, MA and M.D.Shah. 2018. Influence of varying levels of Zn on quality attributes and shelf life of onion in Kashmir. The Parma Innovation Journal, 7(7): 342-345**
- Vijay Aske, P.K. Jain, Narayan Lal and Govind Shiurkar.2017. Effect of Micronutrients on Yield, Quality and Storability of Onion cv. Bhima Super. Trends in Biosciences 10 (6): 1354-1358**
- Walkley AJ and C.A.Black. 1934 An estimation of the digestion method of determining soil organic matter and proposed modification of the chromic acid titration method. Soil Science. 37:29-38.**
- Yoldas F., Ceylan S., Mordogan N. and B.C.Esetlili. 2011. Effect of organic and inorganic fertilizers on yield and mineral content of onion (*Allium cepa* L.). African Journal of Biotechnology, 10: 11488–11492.**

DRAFT