



RESEARCH ARTICLE

**Biochemical Profile of Transgenic Banana cv. Rasthali (*Musa* spp.)
Engineered for Resistance Against *Fusarium* Wilt**

Deepa Sankari R¹, Kumar K K², Varanavasiappan S², Kalpana K¹, Sunisha C⁶,
Usharani T R⁶, Muthulakshmi P⁴, Auxcilia J⁵, Arul L², Kokiladevi E², Nakkeeran
S³, Sudhakar D^{2*} and Eraivan Arutkani Aiyathan K^{1*}

¹Department of Plant Pathology, Agricultural College and Research Institute,
Madurai - 625 104

²Department of Plant Biotechnology, Centre for Plant Molecular Biology and
Biotechnology, Tamil Nadu Agricultural University, Coimbatore-641003

³Agricultural College and Research Institute, Kudumiyanmalai-622 104

⁴Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore-
641003

⁵Department of Fruit Science, Horticultural College and Research Institute, Tamil
Nadu Agricultural University, Coimbatore-641003

⁶Division of Biotechnology, ICAR- Indian Institute of Horticultural Research,
Hesarghatta Lake post, Bangalore - 560 089

ABSTRACT

The biochemical and metabolic profiles of transgenic banana (*Musa* spp.) plants developed through RNAi technology for expressing resistance to *Fusarium oxysporum* f.sp. *cubense* race 1 were characterized by using gas chromatography coupled to electron impact ionization-time of flight-mass spectrometry (GC/EITOF-MS). A total of twenty-two metabolites were identified and annotated using NIST data library. Several resistance related metabolites such as 13-docosenamide, 9-octadecenamide, 9-octadecene, n-hexadecanoic acid, oleic acid, 1,2-benzenedicarboxylic acid, 2,4,6-decatricenoic acid, 1 bis (2-ethylhexyl) phthalate, methyl abietate, undecanedioic acid, 2-pentadecanone, methyl palmitate and phenols were detected in transgenic plants which were generated to silence two fungal genes viz., *ftf1* and/or *velvet* through RNAi technology. The induction of defense related enzyme activities such as phenylalanine ammonia lyase (PAL), superoxide dismutase (SOD), peroxidase (POD) and polyphenol oxidase (PPO) were also observed. The defense related enzyme activities were higher in transformed plants than untransformed plants. However,

these differences did not cause any unintended changes in the growth of transgenic banana plant.

Keywords: *Transgenic banana; Ftf1 gene; Velvet gene; RNAi; Fusarium oxysporum f.sp. cubense race 1; GC/EITOF-MS*

INTRODUCTION

Fusarium wilt of banana, popularly known as Panama disease is a serious fungal disease caused by the soil-borne fungus *Fusarium oxysporum* f.sp. *cubense* (*Foc*) that significantly affects banana production in India. The fungus enters the plant through the roots, colonizing the xylem vessels and thereby blocking water and nutrients flow. Disease progression results in the collapse of leaves at the petiole, the splitting of the pseudostem base and eventually plant death (Ploetz, 2015; Dita *et al.*, 2018). Different races of the pathogen were identified based on the pathogenicity to reference host cultivars: race 1 (R1) affects Gros Michel (AAA) and Manzano/Apple/Latundan (Silk, AAB); race 2 (R2) affects cooking bananas of the Bluggoe (ABB) subgroup; race 3 (R3) affects *Heliconia* spp., and race 4 (R4) affects all cultivars in the Cavendish (AAA) subgroup in addition to those susceptible to R1 and R2 (Waite and Stover, 1960). Currently, all commercial banana cultivars in India are susceptible to *Fusarium* wilt, and it is difficult to manage the disease using chemical fungicides. A sustainable solution for producing bananas in infected soils is replacing susceptible cultivars with resistant ones. Due to lack of suitable *Fusarium* wilt resistant resources, transgenic resistance by exploiting the concept of post transcriptional gene silencing (PTGS) has been employed to manage the disease (Bag *et al.*, 2007; Mehta *et al.*, 2013).

Metabolomics as a biochemical tool in systems biology has the potential application not only for gene function elucidation, but also as a tool to screen for disease resistance. As with genomics, transcriptomics, and proteomics studies, metabolomics generates large data sets of metabolites (Razzaq *et al.*, 2019). Identifying a set of metabolites could explain the functions associated with a trait or gene. Plant extracts usually comprise many secondary metabolites like terpenoids, saponins, alkaloids phenolics flavonoids and glycosides, which possess strong antifungal effects. Several higher plants and their constituents have been reported to control a number of pathogens that cause severe plant diseases (Elzaki *et al.*, 2012; Munawwar *et al.*, 2018; Khan *et al.*, 2018; Bashir *et al.*, 2019 and Hanif *et al.*, 2017). The identification of metabolites is very important for understanding the physiology of plants, including plant growth, development, and defense mechanisms which have been elucidated clearly in transgenic *Nicotiana tabacum* (Lucas *et al.*, 2003, 2006) and transgenic groundnut (Shomo *et al.*, 2016) against viral and fungal diseases. Research evidence showed that the level of expression of phenylalanine ammonia lyase (PAL3) and phenol oxidising enzymes increased in cavendish Brazilian banana when infected with *Foc* TR4 strain compared with non-virulent *Foc* strain TR1 (Chittor *et al.*, 1999; De Ascensao and Dubery, 2000 and Li *et al.*, 2013). Employing RNAi technology, transgenic banana lines (cv. Rasthali) harboring two different RNAi constructs targeting *fusarium transcription factor1* (*ftf1*) and *velvet* gene of *Foc* were generated (unpublished). Analysis of these transgenic lines for *Foc* 1 resistance revealed that transgenic lines expressed better resistance than the untransformed control lines. In the present study, the metabolic profiling of transgenic banana plants was compared with that of un-transformed control by using gas chromatography coupled to electron impact ionization time of flight mass spectrometry (GC/EI-TOF-MS). Besides, the activity of stress related enzymes was also determined in transgenic banana to understand if there were any significant differences in the biochemical profile of transgenic banana compared to their non-transgenic counterpart.

MATERIAL AND METHODS

Plant material

Ten transgenic Rasthali plants (harboring RNAi construct of *ftf1* or *ftf1+velvet* RNAi), developed in our lab were used in this study. These plants were generated through *Agrobacterium* mediated transformation using RNAi constructs with a view to silence fungal genes, *fusarium transcription factor 1* (*ftf1*) or *ftf1* and *velvet* genes. These plants were found to exhibit resistance against *Fusarium oxysporum* f.sp. *cubense* (*Foc1*) race 1 (6 months post inoculation) in an earlier study (results not published). Five

plants in each of these constructs were selected based on disease severity score and PCR assays. These plants were grown in the transgenic greenhouse under controlled conditions along with untransformed plants.

Fungal strain and virulence assay

Highly virulent *Foc* race 1 (strain KP) was used in this study. Mycelial disc of *Foc* race 1 was multiplied in PDA medium. A nine mm disc of *Foc* race 1 was inoculated into sterile Potato Dextrose Broth and incubated in orbital rotary shaker at 28 ± 2 °C (150 rpm) for five days. Subsequently, the suspension was homogenized, and the conidial load was adjusted to 10^6 colony forming unit (cfu) mL^{-1} . This suspension was used to challenge both transgenic and non-transgenic plants. Six months old *Fusarium* resistant transgenic Rasthali plants together with untransformed plants were used for challenge inoculation with *Foc* 1. Prior to inoculation, plants were removed from pots; roots were trimmed to a length of ~40 cm and were individually inoculated by dipping the roots for 30 min in the conidial suspension comprising of 10^6 conidia mL^{-1} . The plants were incubated for 28 days in a transgenic greenhouse maintained at 28 °C. The disease symptoms were recorded at 0 h, 24 h, 48 h, 72 h and 28 days of post-inoculation. For resistance evaluation, the standard evaluation method of Perez-Vicente *et al.* (2014) was adopted.

Sample preparation for metabolite assay using GC/EITOF-MS

Root samples (young roots or newly emerged roots) from the resistant transgenic plants and untransformed plants (cv. Rasthali) infected with *Foc* race 1 were collected after 72 h of inoculation and stored at -80 °C until metabolite extraction. Polar primary metabolites were extracted from the root tissues and then subjected to derivatization by following the procedure described by Erban *et al.* (2007). Root samples were ground to a fine powder using a pre-chilled mortar and pestle. The root powder (120 mg) was transferred to 2 mL round bottom microcentrifuge tubes, and 1.4 mL of 100 % methanol and 60 μL of internal standard ribitol were added and mixed well by vortexing. The mixtures were incubated at 70 °C for 10 min on a rotary orbital shaker at 150 rpm and then centrifuged at $11,000 \times g$ for 10 min. To the supernatant, 750 μL of chloroform and 1.4 mL of double distilled water were added and vortexed for uniform mixing. The mixture was centrifuged at $2,200 \times g$ for 15 min and 1 mL of the supernatant was transferred to 1.5 mL of micro centrifuge tube and concentrated in a vacuum concentrator. The dried samples were then stored at -70 °C until derivatization. For derivatization, 300 μL of methoxy amination reagent (5 mg of 4- dimethyl aminopyridine and 40 mg of methoxy amine hydrochloride dissolved in 1 mL of pyridine) was added to the micro centrifuge tube and shaken at 37 °C for 2 h. The sample was vortexed every 30 min during 2 h of shaking to dissolve the sample completely in the reagent. After methoxy-amination, the samples were centrifuged at $10,000 \times g$ for 1 min and 40 μL of the supernatant was transferred to 1.5 mL microcentrifuge tube and 70 μL of N, O-Bis (trimethylsilyl) trifluoro acetamide (BSTFA) and 10 μL of retention time standards (n-alkanes) were added to the sample and shaken at 37 °C for 30 min. The n-alkane mixture was prepared by using pyridine at a final concentration of 0.22 mg mL^{-1} each. The samples were then centrifuged at $10,000 \times g$ for 1 min and 95 μL of the supernatant was transferred to GC-MS glass vial for analysis.

Gas chromatography and mass spectrometry (GC-MS) analysis

The sample was injected into column Elite5MS (100% Dimethyl poly siloxane), 30 x 0.25 mm x 0.25 μm equipped with GC Clarus 500 (Perkin Elmer). The turbo Mass-Gold-Perkin-Elmer detector was used. The carrier gas flow rate was 1 mL min^{-1} , split 10:1, and injected volumes were 3 μL . The column temperature was maintained initially at 110 °C at the rate of $10 \text{ }^\circ\text{C min}^{-1}$ no hold followed by an increase up to 280 °C at the rate of $5 \text{ }^\circ\text{C/min}$ to 9 min (hold). The injector temperature was 250 °C, which was held constant for 36 min. The electron impact energy was 70 eV, jule line temperature was set at 2000 °C and the source temperature was set at 200 °C. Electron impact (EI) mass scan (m/z) was recorded in the 45-450 amu (atomic mass unit) range. An ion mass spectrometer and OMA detector were used to monitor the eluted compounds from 10 to 30 min (total analysis time 35min).

Data processing and identification of metabolites

The unprocessed raw data from the GC/MS was converted to a readable format by AMDIS (Automated Mass Spectral Deconvolution and Identification System Version 2.64). The AMDIS software was linked to the NIST (National Institute of Standards and Technology) MS library (Version 06) (Kopka *et al.*, 2005; Erban *et al.*, 2007) to determine the best mass spectral matches and the poor matches (<60%) were considered unidentified. The output data from GC/MS scans, the mass ions and their areas were processed by running the AMDIS software. Each metabolite across the replicates was manually verified for consistency using retention time as a reference. Compound structures were putatively identified and evaluated by comparing the molecular masses (m/z values) of the eluted compounds with literature data and standards.

Assessment of defense related enzymes activity

The samples from the most recent completely expanded leaf of the infected plants were used for biochemical analysis. The phenylalanine ammonia lyase enzyme activity was carried out at 24 h intervals after *Foc1* inoculation. Meanwhile, peroxidase, polyphenol oxidase and superoxide dismutase activities were measured at 0 h and 28 days after inoculation. All biochemical parameters were measured using a spectrophotometer (Jasco V-730 BIO spectrophotometer, USA).

Phenylalanine ammonia lyase assay (PAL)

PAL was assayed according to the method of Robert and Helmut (1992). The absorbances of the reaction mixture of the samples were measured at 290 nm with its corresponding control at every 30 min intervals. PAL activity was expressed as changes in absorbance at 290 nm h⁻¹ mg⁻¹ protein.

Peroxidase assay

Peroxidase activity was assessed following the oxidation of pyrogallol according to the method given by Hammerschmidt *et al.* (1982). The absorbance was recorded at 430nm at 30 sec intervals up to 3 min. Peroxidase activity was expressed in unit mg⁻¹ protein. A change in 0.1 absorbance min⁻¹ mg⁻¹ protein refers as one unit.

Polyphenol oxidase assay

Polyphenol oxidase activity was determined according to the method described by Mozzetti *et al.* (1995). The activity was measured by monitoring the increase in absorbance for 3 min at 410 nm. The specific activity was expressed in unit mg⁻¹ protein. Similar to peroxidase, a change in 0.1 absorbance min⁻¹ mg⁻¹ protein refers as one unit.

Superoxide Dismutase (SOD)

The SOD activity was estimated by following the method described by Du and Bramlage (1994). The inhibition of photochemical reduction of nitro blue tetrazolium (NBT) was measured at 560 nm and expressed in units g⁻¹ fresh weight. The amount of enzyme that inhibited the NBT photo reduction by 50% was accounted as one unit of SOD activity.

Statistical analysis

Data were analyzed using one-way ANOVA and the differences contrasted using Duncan's multiple range test. All statistical analyses were performed at the level 5% using SPSS 10.0 (SPSS Inc.,).

RESULTS AND DISCUSSION

Bioassay of transgenic banana expressing *ftf1* and *ftf1* + *velvet* gene sequences

The selected *Foc* 1 resistant transgenic banana plants derived from the two RNAi constructs were subjected to *Foc* race 1 infection. *Fusarium* wilt symptoms were observed in untransformed control banana plants only after 28 days of inoculation. Infected plants were characterized with yellowing of older leaves coupled with splitting of the pseudostem. Subsequently, infected plants wilted completely after 6 weeks. However, there was no visible symptom on non-inoculated transgenic plants after 28 days of post inoculation. Subsequently, few transgenic plants (FTF-289, 335; FTF+VELVET (F+V) - 51, 116) exhibited yellowing of older leaves and pseudostem splitting, whereas the plants viz., FTF-215; FTF+VELVET-93, 97 were free of *Foc* symptoms at 28 days post inoculation (DPI) (Figure 1).

Figure 1. Bioassay of the transgenic banana plants for *Foc* resistance at 28 days post inoculation

a) UC 1- Rasthali control (non-inoculated), UC 2- Rasthali control (inoculated), UC-3 Grand naine control (inoculated), *ftf1* plants and b) UC 1- Rasthali control (non-inoculated), UC 2- Rasthali control (inoculated), UC-3 Grand naine control (inoculated), *ftf1+velvet* plants (UC-untransformed control)

Results from the disease incidence as shown in Figure 2, indicated that the disease incidence among transgenic plants varied between 10 and 80% from both the construct derived plants. Ghag *et al.* (2014) reported that the differences in disease resistance of Rasthali transgenic plants with wild type against *Foc* in banana could depend on genotype, environmental conditions and virulence of the fungal strain.




Figure 2. Disease incidence and disease grade of transformed Rasthali plants against *Foc* race 1 (28 DPI). Data were analyzed using one-way ANOVA and the differences contrasted using Duncan's multiple range test. Values indicated in superscripts (a-e) are significantly different ($p < 0.05$)

Metabolite compounds in transgenic banana

Fungal resistance in a variety of crop plants has been successfully achieved by transforming the host plant with genes derived from fungus. The entire or partial gene sequences of the pathogen (both structural and functional) are used to transform host plants to generate dsRNA in order to protect them against the pathogen or a range of closely related pathogens (Baulcombe, 1996). This strategy has been employed to protect crop plants against several genera of fungus in various plant species (Galvez *et al.*, 2014). Detailed analysis of transgenic plants revealed that resistance in plants is primarily due to transgene induced post-transcriptional gene silencing (PTGS) via the formation of dsRNA, termed as "RNA silencing" (Baulcombe, 2002). In the present study, the metabolic profile of transgenic Rasthali banana plants transformed with gene construct encoding dsRNA *ftf1* and/or *velvet* gene sequence were compared with untransformed counterparts against *Foc* race 1.

The samples of resistant transgenic plants (FTF-215, FTF1+VELVET-97) and susceptible wild type plant after 72 h of inoculation with *Foc* race 1 were chosen for metabolite analysis based on the disease severity score obtained 28 DPI. The parameters such as retention time, molecular formula, area percentage, and biological activity of different compounds were identified by the comparison of their mass spectra against the NIST library database (Table 1). The peak area represented a quantitative proportion of the predicted compound to the total of crude extracts. These compounds with antifungal actions were related to various chemical classes, including esters, fatty acids, aldehydes, tertiary amines, alkaloids, and ketones. Among these bioactive compounds obtained, transgenic plants FTF-215 and F+V-97 had significant area (per cent) in comparison with control plants, indicating that they played a substantial role in antifungal activities. Results obtained on the GCMS analysis of the transgenic and untransformed banana plants are presented in Figure 3 a, b, c.

Several resistance-related metabolites were consistently detected in both transgenic and untransformed plants. The differentially expressed bioactive metabolites during the interaction of transgenic plants with *Foc1* KP revealed that transgenic plants produced seven unique bioactive metabolites such as 1,2-benzenedicarboxylic acid, undecanedioic acid, 2-pentadecanone, bis (2-ethylhexyl) phthalate, methyl palmitate, methyl abietate, and 2,4,6-decatricenoic acid exclusively in the transgenic plant (Table 1).

Likewise, Rasthali wildtype with *Foc* KP infection induced secretion of secondary metabolites *viz.*, octadecanoic acid methyl ester, eicosane, heptadecane, 2,6-dimethyl, 4,6-dimethyldodecane, trioxsalen, hexanoic acid and butanamide. Comparison of bioactive metabolites between Rasthali wild type with *Foc* KP and *ftf1* or *ftf1+velvet* with *Foc* KP revealed the production of 15 bioactive metabolites in common *viz.*, 9-octadecenoic acid, octadecane, 2-ethyl-3-methylpyrazine, 2-methyl-5,6 diethyl pyrazine, phenol 2,4-bis (1,1-dimethylethyl)-4-methyl, octadecanoic acid, 1-ascorbic acid, n-hexadecanoic acid, 1-butanone, 9-octadecenamide, ethyl iso-allocholate, 9-octadecene, lauric acid and oleic acid (Figure 3 a, b, c). In this study, a volatile resin, 2-pentadecanone detected exclusively in transgenic plants exhibiting a higher peak area percentage of 8.92, thereby indicating that this metabolite plays a potential role in plant defense by exhibiting antifungal activity similar to other fungal pathogens such as *Aspergillus* sp., *Fusarium oxysporum*, *Pythium* sp., and *R. solani* in beans (Hanif *et al.*, 2022).

The metabolite, 13-docosenamide or erucamide was detected at a retention time of 25.45 min with 2.66 % area increase in transgenic plants. Generally, it acts as anti-inflammatory, dermatogenic, hypocholesterolemic and anaemiagenic (Li *et al.*, 2017). The antifungal mechanism of erucamide, destroy cell structure and inhibit the germination and growth of fungal spores (Makarieva *et al.*, 2002). Enhanced disease resistance in plants induced by erucamide has been demonstrated against *Foc* TR4 in banana (Qi *et al.*, 2019). The metabolite, 9-octadecenamide, expressed in transgenic Rasthali expressing siRNA specific to *ftf1* and *velvet* has been shown to have antifungal activity against *Botrytis cineria* and *Macrophomina Phaseolina* (Khan *et al.*, 2022) in soybean. Dodecanoic acid (Lauric acid), a saturated fatty acid was detected at a retention time of 28.52 min in transgenic plants (2.25 %), and is known to exhibit antifungal activity against *Foc*. It also exhibits antimicrobial activity against various bacteria and antifungal

activity against *Pythium ultimum*, *Rhizoctonia solani*, *Blumeria*, *Aspergillus* and *Fusarium* spp. (Walters et al., 2003; Liu et al., 2008).

The metabolite, 2-methyl-5,6 diethyl pyrazine, which occurs in much larger amounts in bacteria and in some plants was detected at a retention time of 9.25 in transgenic plants with an area percentage of 6.05 (*ftf1*) and 6.25 (*ftf1+velvet*). The antifungal activity of the metabolite was reported earlier against *Fusarium solani*, *Fusarium* sp., and *Colletotrichum gloeosporioides* (Edgar et al., 2019). Rahman and Anwar (2006) reported the broad-spectrum antifungal activity of 1,2-benzenedicarboxylic acid against six phytopathogenic fungi viz., *Alternaria alternata*, *Botryodiplodia theobromae*, *Curvularia lunata*, *Fusarium equiseti*, *Macrophomina phaseolina* and *Colletotrichum corchori*. Similarly, the metabolite bis (2-ethylhexyl) phthalate exhibited antimicrobial, antioxidant and antifungal activities against unicellular and filamentous fungi like *Fusarium oxysporum* (Zishan et al., 2020). The antifungal activity of 1-ascorbic acid which is produced by transgenic plants was reported by Bokhari et al. (2013) against *Fusarium oxysporum*, *Alternaria alternata*, *Macrophomina phaseolina* and *Colletotrichum* in *Citrullus* sp. Shobier et al. (2016) reported the antibacterial and antifungal activities (*Aspergillus* spp., *Fusarium* sp.,) of methyl palmitate that damages microbial cellular membranes in various crops. Likewise, methyl abietate possessing high antioxidant activity, exhibited antibacterial activity against *E. coli*, *B. cereus* and antifungal activity against *Puccinia* sp., *Fusarium oxysporum*, *Alternaria alternata* and *Fusarium gramineum* in several crops.



Table 1. Metabolite profiles of transgenic and wild type banana plants assessed by GC-MS upon inoculation with *Foc* race 1

S. No.	Retention time (min)	Name of the compound with molecular formula	Peak area (%) in wild type plant	Peak area (%) in transgenic plant		Inhibitory role / Biological activity	Reference
				FTF1	F+V		
1	3.02	9-Octadecenoic acid (Ricinoleic acid), C ₂₁ H ₄₀ O ₄	7.56	9.74	9.84	Antibacterial activity against <i>E. coli</i> , <i>Salmonella</i> sp. and antifungal activity against <i>Aspergillus flavus</i> and <i>Fusarium</i> spp.	Krishnaveni et al. (2014)
2	4.64	Octadecane, C ₄₂ H ₈₆	0.76	1.12	1.15	Antifungal activity against <i>Fusarium</i> sp. and <i>C. gloeosporioides</i>	Edgar et al. (2019)
3	6.41	2-Ethyl-3-methylpyrazine, C ₇ H ₁₀ N ₂	2.12	4.02	3.99	Inhibits the growth of <i>Fusarium oxysporum</i> , <i>Botryosphaeria</i> sp., <i>Trichoderma atroviride</i> , <i>Colletotrichum gloeosporioides</i> and <i>Penicillium expansum</i>	He et al. (2020)
4	9.25	2-Methyl-5,6 diethyl pyrazine, C ₉ H ₁₄ N ₂	1.97	6.05	6.25	Antifungal activity against <i>Fusarium solani</i> , <i>Fusarium</i> sp. and <i>Colletotrichum gloeosporioides</i>	Edgar et al. (2019)
5	9.64	1,2-Benzenedicarboxylic acid, C ₈ H ₆ O ₄	-	1.67	1.71	Antifungal activity against six phytopathogenic fungi viz., <i>Alternaria alternata</i> , <i>Botryodiplodia theobromae</i> , <i>Curvularia lunata</i> , <i>Fusarium equiseti</i> , <i>Macrophomina phaseolina</i> and <i>Colletotrichum corchori</i> .	Rahman and Anwar (2006)
6	10.95	Undecanedioic acid, C ₁₁ H ₂₂ O ₂	-	5.74	5.87	Antifungal activity against <i>Fusarium oxysporum</i> f. sp. <i>cubense</i>	Yang et al. (2021)
7	12.85	2-Pentadecanone, C ₁₅ H ₃₀ O	-	8.96	8.88	Antifungal activity against <i>Aspergillus</i> sp., <i>Fusarium oxysporum</i> , <i>Pythium</i> sp. and <i>Rhizoctonia solani</i>	Hanif et al. (2022)
8	14.18	Phenol 2,4-bis (1,1-dimethylethyl)-4-methyl, C ₁₄ H ₂₂ O ₂	0.33	3.92	3.99	Completely inhibited the growth of <i>Fusarium oxysporum</i> f.sp. <i>cubense</i>	Yuan et al. (2012)
9	18.21	Octadecanoic acid (Stearic acid), C ₁₈ H ₃₆ O ₂	0.36	1.47	1.45	Exhibit antifungal property against <i>F. oxysporum</i> and <i>M. phaseolina</i>	Aftab et al. (2019)
10	20.36	Bis (2-ethylhexyl) phthalate, C ₂₄ H ₃₈ O ₄	-	1.8	1.9	Exhibited antimicrobial, antioxidant and antifungal activities against unicellular and filamentous fungi like <i>Fusarium oxysporum</i>	khadim et al. (2016)
11	22.21	L-Ascorbic acid, C ₃₈ H ₆₈ O ₈	1.69	5.77	5.89	Antifungal activity against <i>Fusarium oxysporum</i> , <i>Alternaria alternata</i> , <i>Macrophomina phaseolina</i> and <i>Colletotrichum musae</i>	Bokhari et al. (2013)

12	22.21	n-Hexadecanoic acid (Palmitic acid), C ₁₆ H ₃₂ O ₂	1.77	9.20	9.34	Exhibited antifungal properties against a number of fungal species namely <i>Fusarium sporotrichioides</i> , <i>F. subglutinans</i> , <i>F. verticillioides</i> , <i>F. cerealis</i> , <i>F. proliferatum</i> , <i>F. oxysporum</i> , <i>Aspergillus carbonarius</i> , <i>A. tubingensis</i> , <i>Penicillium</i> sp. and <i>Alternaria alternata</i>	Morcia et al. (2012)
13	24.24	Methyl palmitate, C ₁₇ H ₃₄ O ₂	-	2.21	2.30	Exhibited antibacterial and antifungal activities (<i>Aspergillus</i> spp., <i>Fusarium</i> sp.) by damaging microbial cellular membranes	Shobier et al. (2016)
14	24.27	1-Butanone, C ₄ H ₈ O	0.56	0.89	0.97	Inhibits the growth of <i>Fusarium</i> spp.	Khan and Javaid (2022)
15	25.45	13-Docosenamide (Erucamide), C ₂₂ H ₄₃ NO	0.57	3.19	3.23	Antibacterial and antifungal activity by destroying the cell structure and inhibit the germination and growth of fungal spores in <i>Foc</i> TR4	Qi et al. (2019)
16	25.57	9-Octadecenamide, C ₁₈ H ₃₅ NO	1.07	3.19	3.17	Exhibits antifungal activity against <i>Colletotrichum acutatum</i> , <i>C. coccodes</i> , <i>C. gloeosporioides</i> , <i>F. oxysporum</i> , and <i>T. roseum</i> .	Kim et al. (2019)
17	27.21	Ethyl iso-allocholate, C ₂₆ H ₄₄ O ₅	0.53	1.96	1.99	High inhibitory activity against different strains of gram-negative bacteria, anti-diabetic activity and resistance property against <i>Fusarium oxysporum</i> f.sp. <i>lycopersici</i>	Rodriguez et al. (2015)
18	27.85	Methyl abietate, C ₂₁ H ₃₂ O ₂	-	1.95	1.96	Antibacterial against <i>E. coli</i> , <i>B. cereus</i> , antifungal activity against <i>Puccinia</i> sp., <i>Fusarium oxysporum</i> , <i>Alternaria alternata</i> , <i>Fusarium gramineum</i> and high antioxidant activity	Benlembarek et al. (2021)
19	28.17	2,4,6-Decatrienoic acid, C ₁₀ H ₁₄ O ₂	-	2.23	2.20	Antimicrobial activity against five bacteria and fourteen fungi (includes <i>Fusarium</i> spp.) and yeast.	Kadhim et al. (2016)
20	28.48	9-Octadecene, C ₃₈ H ₇₆ O ₂	0.34	2.27	2.28	Antifungal, antioxidant, anticarcinogenic, and antimicrobial activity against yeast fungus, <i>S. rolfisii</i>	Akpuaka et al. (2013)
21	28.52	Dodecanoic acid (Lauric acid), C ₁₂ H ₂₄ O ₂	1.22	2.27	2.26	Antimicrobial activity against various bacteria and antifungal activity against <i>Pythium ultimum</i> , <i>R. solani</i> , <i>Blumeria</i> , <i>Aspergillus</i> and <i>Fusarium</i> spp.	Liu et al. (2008)
22	29.06	Oleic acid, C ₁₈ H ₃₄ O ₂	0.32	2.27	2.35	Antifungal activities against <i>Aspergillus niger</i> , <i>A. flavus</i> , <i>Penicillium parasiticus</i> , <i>Candida utilis</i> , <i>Fusarium solani</i> and antibacterial activity against gram-positive bacteria.	Mohy El-Din and Mohyeldin (2018)




Figure 3. Total ion chromatogram of transgenic plants at 72 h post inoculation (A) FTF-215 (B) FTF+VELVET-97 and (C) wild-type banana plants

The metabolite, undecanedioic acid was detected at a retention time of 10.95 min in transgenic plants alone with an area percentage of 5.74 (*ftf1*) and 5.87 (*ftf1+velvet*). Antifungal activity of undecanedioic acid in plants has been reported earlier against *Fusarium oxysporum* f. sp. *cubense* in banana by Yang *et al.* (2021). A phenolic compound, phenol, 2,6-bis (1,1-dimethylethyl)-4-methyl, commonly known as butylated hydroxytoluene is an antioxidant. This metabolite which has been demonstrated as an antimicrobial agent due to its ability to remove free radicals was also detected in transgenics (Pintac *et al.*, 2019). Yuan *et al.* (2012) reported that the presence of phenol completely inhibited the growth of *Fusarium oxysporum* f. sp. *cubense* in transgenic banana.

The metabolite, 9-octadecenoic acid or ricinoleic acid was detected in the transgenic plant and wild type plant at a retention time of 3.02. The ricinoleic acid area percentage in wild type was 7.56 %, whereas 9.74 % and 9.84 % were recorded in *ftf1* or *ftf1+velvet* plants, respectively. The concentration of ricinoleic acid increased as infection progressed and was implicated in the activation of defense responses against injury/infection of *Aspergillus flavus* and *Fusarium* spp. (Krishnaveni *et al.*, 2014). Octadecanoic acid or stearic acid was detected at a retention time of 18.21 min with an increase in 1.11 % area in transgenic plants compared to wild type. Stearic acid is known to exhibit antifungal property against soil borne diseases such as *F. oxysporum* and *M. Phaseolina* in black cumin (Aftab *et al.*, 2019).

In this study, the differences in peak area percentage of various metabolites among transgenics and wild type were compared. However, these differences did not cause any changes in the growth and development of transgenic plants. Besides, similar observations on the production of metabolites in Tobacco streak virus TSV) resistant transgenic groundnut plants engineered to express dsRNA of coat protein gene of TSV (Shomo *et al.*, 2016) and transgenic cotton plants engineered for down-regulation of tryptophan synthesis-related gene (*GbTRP1*) for controlling *Verticillium dahliae* and *Botrytis cinerea* (Miao *et al.*, 2019).

Phenylalanine ammonia lyase activity in transgenic plants

L-Phenylalanine ammonia lyase, in phenylpropanoid pathway acts as an inducible physical barrier against pathogen ingress and has shown to be strongly induced with the infection of fungal pathogen, *Cercospora nicotianae* in transgenic tobacco (Gail *et al.*, 2003). In this study, PAL activity was monitored in Rasthali (AAB) transformed and untransformed plants following inoculation with *Foc* race 1 up to 72 h. PAL activity increased to maximum level by 24 h in transgenic banana plants (F+V 97; FTF 215) that correlates to the disease incidence obtained at 28 DPI with *Foc1* (Table2; Figure 2). Similar observation was reported by Aguilar *et al.* (1999) in gold finger cultivar (AAAA), which is known to be more resistant to *Fusarium* wilt disease with the accumulation of lignin (De Ascensao and Dubery 2000) when compared with banana cultivar susceptible to *Fusarium* wilt such as Williams (AAA) and Gros Michel (AAA). The observation of the present study is also supported by the findings of Subramaniam *et al.* (2006), who showed that the transgenic banana Rasthali-pisang cultivar (AAB) expressing two genes (*chitinase* and β -1,3 *glucanase*) had higher PAL activity against *Foc* after 24 h of inoculation.

Peroxidase, superoxide dismutase and polyphenol oxidase activity in transgenic banana plants during pre and post inoculation with *Foc*

In stress-related environment, ROS accumulation appears to be at a higher level and found to be a common phenomenon. The activity of ROS scavenging enzymes viz., SOD, POD and PPO were studied in transgenic plants before and in response to pathogen infection. SOD detoxifies the O_2^- radicals to H_2O_2 and acts as the first line of defense mechanism, POD detoxifies the peroxides into H_2O and serves as the next line of defense by formation of lignin or suberin, a physical barrier to limit pathogen invasion and enhance cell wall rigidity. Further, PPO is responsible for oxidation of phenolic compounds into antimicrobial quinones in plant cells attacked by phytopathogens (Malolepsza and Urbanek, 2000). In the present experiment, the activity of these ROS enzymes substantially increased after 28 days of *Foc* inoculation in

the transgenic plants compared to the untransformed ones. The results obtained also correlates to the disease severity score.

Table 2. Time course of changes in Phenylalanine ammonia lyase (PAL) activity (*Foc 1* post inoculation) in control and transgenic Rasthali (AAB) plants


Plant Id	Hours			
	0	24	48	72
Rasthali control	3.7 ^a	12.1 ^b	22.5 ^{cd}	6.4 ^a
Grand Naine control	4 ^a	16.9 ^c	25.3 ^d	13.6 ^b
FTF 160	4.3 ^a	16.5 ^c	25.1 ^d	14.5 ^b
FTF 215	4.9 ^a	20.6 ^c	26.9 ^d	15.8 ^c
FTF 289	3.6 ^a	14.5 ^b	21.2 ^c	11.9 ^b
FTF 335	3.9 ^a	15.3 ^c	23.4 ^d	13.7 ^b
FTF 428	4.1 ^a	16.4 ^c	24.5 ^d	14.3 ^b
F+V 51	4.6 ^a	15.7 ^c	19.6 ^c	11.4 ^b
F+V 93	5.2 ^a	19.3 ^c	25.6 ^d	15.1 ^c
F+V 97	6.2 ^a	21.1 ^c	27.8 ^d	16.9 ^c
F+V 116	5.9 ^a	17.2 ^c	22.3 ^{cd}	13.1 ^b
F+V 120	5.4 ^a	19.1 ^c	24.7 ^d	14.9 ^b

Data were analyzed using one-way ANOVA and the differences contrasted using Duncan's multiple range test. Values indicated in superscripts (a-d) are significantly different ($p < 0.05$)


The specific activity of peroxidase was higher (46%) in the transgenic plant F+V-97 (54.3-unit mg^{-1} protein) without disease symptom expression when compared to the untransformed plant (24.8-unit mg^{-1} protein) (Figure 4a) suggesting that peroxidase enzyme could be involved in defense response against pathogens. Morpurgo *et al.* (1994) reported that the activity of peroxidase was five times higher in the leaves, roots, and corm tissues of the *F. oxysporum* resistant banana cultivar, Gold Finger (AAAA) than in the susceptible cultivar, Pisang Mas (AA). Subsequently, the increase in superoxide dismutase activity was recorded as 47 and 46.5 per cent higher in transgenic plants F+V-97, FTF-215 (58.3, 57.5 unit mg^{-1} protein) compared to untransformed plant (26.8 unit mg^{-1} protein) (Figure 4b). Similar observations were made by Sunisha *et al.* (2020) after stacking of antimicrobial genes in banana against *Fusarium* wilt and Karmakar (2017) in the leaf sheath of transgenic lines of rice to *R. solani* infection and reported increased peroxidase, superoxide dismutase and ascorbate peroxidase activities.

Besides, an increase in polyphenol oxidase activity in the transformed Rasthali (AAB) plant was also observed (Figure 4c). The highest PPO activity in F+V 97 (65.09-unit mg^{-1} protein) transgenic plant was recorded, which was 44% higher than that recorded from untransformed plants. Subsequently, other transformed plants *viz.*, F+V-93 and FTF-215, it varies significantly from each other, where the PPO activity ranged between 56.3 and 62.3 unit mg^{-1} protein. Similar findings were observed by Mohammadi and Kazemi (2002) and Subramaniam *et al.* (2006) with increased PPO activity in transgenic wheat heads and pisang-rasthali cultivars, respectively, following inoculation with respective *Fusarium* conidia.

a) Peroxidase (POD) activity



b) Superoxide dismutase (SOD) activity



c) Polyphenol oxidase (PPO) activity




Figure 4. Changes in enzymatic activities in wild type and transgenic Rasthali (AAB) plants before and 28 DPI of *Foc*. Data were analyzed using one-way ANOVA and the differences contrasted using Duncan's multiple range test. Values indicated in superscripts (a-d) are significantly different ($p < 0.05$)

The plant defense mechanism is a complex one with the emergence of new pathogenic strains that makes it difficult to study. Considering the pathogenic nature, various biochemical parameters against *Foc* transformed and untransformed banana cv. Rasthali indicated a significant increase in defense enzyme activities in the transformed plants compared to untransformed ones. An increase in the amount of hydrogen peroxide, PAL activities after 24 h, and phenol oxidizing enzymes after 28 days of *Foc* inoculation was found higher in transformed Rasthali (AAB) plants with combined *ftf1+velvet* gene than the single inserted *ftf1* gene.

CONCLUSION

The present study revealed the resistance activity of transgenic banana plants against *Foc* race 1 by understanding their biochemical profile. The bioactive metabolite compounds identified using GC-MS analysis support and signifying the importance of the resistant action of transgenic plants against *Foc*. However, detailed characterization of these antifungal compounds needs to be carried out to know the exact mode of action.

Funding and Acknowledgment

The authors would like to thank DBT-NER BANANA (GOI-DBT/NER BANANA/2017-18) for the funding and Centre for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University, Coimbatore for providing facilities. RDS also thank DBT-NER banana for the fellowship.

REFERENCES

- Aftab, A., Yousaf, Z., Javaid, A., Riaz, N., Younas, A., Rashid, M., Shamsher, B. and A. Arif. 2019. Antifungal activity of vegetative methanolic extracts of *Nigella sativa* against *Fusarium oxysporum* and *Macrophomina phaseolina* and its phytochemical profiling by GC-MS analysis. *Int. J. Agric. Biol.*, **21**: 569-576.
- Aguilar, E. A., Turner, D. W. and K. Sivasithamparam. 1999. Aerenchyma formation in roots of four banana (*Musa* spp.) cultivars. *Sci. Horti.*, **80(1-2)**: 57-72.
- Akpuaka, A., Ekwenchi, M. M., Dashak, D. A. and A. Dildar. 2013. Biological activities of characterized isolates of n-hexane extract of *Azadirachta indica* A. Juss (Neem) leaves. *Nat. & sci.*, **11(5)**: 141-147.
- Bag, S., Singh, R. S. and R. K. Jain. 2007. *Agrobacterium* mediated transformation of groundnut with coat protein gene of *Tobacco streak virus*. *Indian J. Virol.* **18**: 65-69.
- Bashir, S., Jabeen, K., Iqbal, S., Javed, S. and A. Naeem. 2019. Lantana camara: Phytochemical analysis & antifungal prospective. *Planta Daninha.*, **37**: 019193526.
- Baulcombe, D. C. 1996. RNA as a target and an initiator of post-transcriptional gene silencing in transgenic plants. *Plant Mol Biol.*, **32(1-2)**:79-88
- Baulcombe, D. 2002. RNA silencing. *Current bio.*, **12(3)**: R82-R84.
- Benlembarek, K., Lograd, T., Ramdani, M. and P. Chalard. 2021. Chemical composition, antibacterial, antifungal and antioxidant activities of *Taxus baccata* essential oil from Algeria. *Biodiversitas J. of Bio. Div.*, **22 (12)**.
- Bokhari, N. A., Kahkashan, P., Iffat, S., Iram, S., Wahibi, M. S. and D. A. Soliman. 2013. Antifungal activity of *Citrullus colocynthis* against *Fusarium oxysporum*, *Alternaria alternata*, *Macrophomina phaseolina* and *Colletotrichum musae*. *J. of Pure and Appl. Microbiology.*, **7(4)**: 2981-2986.
- Chittor, J.M., Leach, J.E. and F. F. Whit. 1999. Induction of peroxidase during defense against pathogen. In: Datta, S.K., Muthukrishnan, (Eds.), Pathogenesis Related Proteins in Plants. *CRC Press*, Boca Raton, FL, pp. 291–325.
- De Ascensao, A.R.D.C.F and I. A. Dubery. 2000. Panama disease: cell wall reinforcement in response to elicitors from *Fusarium oxysporum* f.sp. *cubense* race four., *Am. Phytopathol. Soc.* **90**: 1173–1180.
- De Rodríguez, D.J., Trejo-González, F.A., Rodríguez-García, R., Díaz-Jimenez, M.L.V., Sáenz-Galindo, A., Hernández-Castillo, F.D., Villarreal-Quintanilla, J.A. and Peña-Ramos, F.M., 2015. Antifungal activity in vitro of *Rhus muelleri* against *Fusarium oxysporum* f. sp. *lycopersici*. *Industrial Crops and Products*, **75**, pp.150-158.

- Dita, M., Barquero, M., Heck, D., Mizubuti, E. S. and C. P. Staver. 2018. Fusarium wilt of banana: current knowledge on epidemiology and research needs toward sustainable disease management. *Front. Pl. Sci.* **9**:1468.
- Du, Z., and W. J. Bramlage. 1994. Superoxide dismutase activities in senescing apple fruit (*Malus domestica* Borkh.). *J. of Food Sci.*, **59**(3):581–584.
- Edgar Guevara-Avendaño., Bejarano-Bolívar, A. A., Kiel-Martínez, A. L., Ramírez-Vázquez, M., Méndez-Bravo, A., von Wobeser E. A., Sánchez-Rangel, D., Guerrero-Analco, J, A., Eskalen, A. and F. Reverchon. 2019. Avocado rhizobacteria emit volatile organic compounds with antifungal activity against *Fusarium solani*, *Fusarium* sp. associated with Kuroshio shot hole borer, and *Colletotrichum gloeosporioides*. *Microbiol Res.* **219**:74-83.
- Elzaki, O. T., Khider, T.O., Omer, S.F. and S. K. Shomeina. 2012. Environment friendly alkaline pulping of *Albizia lebeck* from Sudan. *Nat Sci* **10**(4): 76-82.
- Erban, A., Schauer, N., Fernie, A. R. and J. Kopka. 2007. Non supervised construction and application of mass spectral and retention time index libraries from time-of-flight gas chromatography-mass spectrometry metabolite profiles. *Methods Mol. Biol.* **358**: 19- 38.
- Gail, L. S., Wesley, S.V., Kenneth, L. K., Chen, F., Lamb, C. and R. A. Dixon. 2003. Phenylpropanoid compounds and disease resistance in transgenic tobacco with altered expression of L-phenylalanine ammonia-lyase. *Phytochemistry* **64**: 153–161.
- Galvez, L. C., Banerjee, J., Pinar, H. and A. Mitra. 2014. Engineered plant virus resistance. *Plant Sci.* **228**: 11-25.
- Ghag, S. B., Shekhawat, U. K. and T. R. Ganapathi. 2014. Host induced post transcriptional hairpin RNA mediated gene silencing of vital fungal genes confers efficient resistance against Fusarium wilt in banana. *Plant Biotechnology journal*, **12**(5): 541-553.
- Ghag, S. B., Shekhawat U. K. and T. R. Ganapathi. 2015. Fusarium wilt of banana: biology, epidemiology and management. *Int. J Pest Management.* **61**: 250–263.
- Hammerschmidt, R., Nuckelus, E.M. and J. Kuc. 1982. Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*. *Physiol. Plant Pathol.* **20**: 61–71
- Hanif, S., Jabeen, K., Akhtar, N. and S. Iqbal. 2022. GC-MS analysis & antifungal activity of *Datura metel* L. against *Rhizoctonia solani* Kuhn. *Anais da Academia Brasileira de Ciências*, **94**.
- He, C.N., Ye, W.Q., Zhu, Y.Y. and W. W. Zhou. 2020. Antifungal activity of volatile organic compounds produced by *Bacillus methylotrophicus* and *Bacillus thuringiensis* against five common spoilage fungi on loquats. *Molecules*, **25**(15), p.3360.
- Kadhim, M. J., Mohammed, G. J. and I. H. Hameed. 2016. *In vitro* antibacterial, antifungal and phytochemical analysis of methanolic extract of fruit *Cassia fistula*. *Ori. J of Chemi*, **32** (3):1329.
- Karmakar, S., Molla, K. A., Das, K., Sarkar, S. N., Swapan, K. and K. Datta. 2017. Dual gene expression cassette is superior than single gene cassette for enhancing sheath blight tolerance in transgenic rice. *Sci. Rep.*, **7**: 7900.
- Khan, I. H. and A. Javaid. 2022. Antifungal activity of n-butanol stem extract of quinoa against *Macrophomina phaseolina*. *Pak. J of Botany*, **54**(4): 1507-1510.
- Kim, Y.J., Kim, J.H. and J. Y. Rho. 2019. Antifungal activities of *Streptomyces blastmyceticus* strain 12-6 against plant pathogenic fungi. *Mycobiology*, **47**(3), pp.329-334.
- Kopka, J., Schauer, N., Krueger, S., Birkemeyer, C., Usadel, B., Bergmueller, E., Dormann, P., Weckwerth, W., Gibon, Y., Stitt, M., Willmitzer, L., Fernie, A. R. and D. Steinhauser. 2005. GMD@CSDB: The Golm Metabolome Database. *Bioinfo.*, **21**: 1635-1638.
- Krishnaveni Marimuthu., Nandhini Nagaraj and Dhanalakshmi Ravi. 2014. GC-MS Analysis of Phytochemicals, Fatty acids and Antimicrobial Potency of Dry Christmas Lima Beans. *Int. J. Pharm. Sci. Rev. Res.*, **27**(2), 09: 63-66.
- Li, C., Shao, J., Wang, Y., Li, W., Guo, D., Yan, B., Xia, Y., and M. Peng, 2013. Analysis of banana transcriptome and global gene expression profiles in banana roots in response to infection by race 1 and tropical race 4 of *Fusarium oxysporum* f. sp. *cubense*. *BMC Gen.*, **14**:851

- Li, M.M., Jiang, Z.E., Song, L.Y., Quan, Z.S. and H. L. Yu. 2017. Antidepressant and anxiolytic-like behavioral effects of erucamide, a bioactive fatty acid amide, involving the hypothalamus–pituitary–adrenal axis in mice. *Neurosci. Lett.*, **640**, pp.6-12.
- Liu, W. W., Wei, M. U., Zhu, B. Y., Du, Y. C. and L. I. U. Feng. 2008a. Antagonistic activities of volatiles from four strains of *Bacillus* spp. and *Paenibacillus* spp. against soil-borne plant pathogens. *Agri. Sci. in China*, **7(9)**: 1104-1114.
- Liu, X., Feng, X. Q., Yang, S., Wang, T. P. and Z. X. SU, Z. 2008. Effects of molecular weight and concentration of chitosan on antifungal activity against *Aspergillus niger*. *Iranian polymer J*, **17(11)**.
- Lucas, W. J., Balachandran, S., Park, J. and S. Wolf. 1996. Plasmodesmal companion cell-mesophyll communication in the control over carbon metabolism and phloem transport: insights gained from viral movement proteins. *J. Exp. Bot.* **47**: 1119-1128.
- Lucas, W. J., Olesinski, A., Hull, R. J., Haudenschild, J. S., Deom, C. M., Beachy, R. N. and S. Wolf. 1993. Influence of the tobacco mosaic virus 30 kDa protein on carbon metabolism and photosynthate partitioning in transgenic tobacco plants. *Planta.*, **190**: 88-96.
- Makarieva, T. N., Santalova, E. A., Gorshkova, I. A., Dmitrenok, A. S., Guzii, A. G., Gorbach, V. I. and V. A. Stonik. 2002. A new cytotoxic fatty acid (5Z, 9Z) 22-methyl 5, 9 tetracosadienoic acid and the sterols from the far Eastern sponge *Geodinella robusta*. *Lipids.*, **37(1)**: 75-80.
- Malolepsza, U. and H. Urbanek. 2000. The oxidants and antioxidants enzymes in tomato leaves treated with o-hydroxyethylrutin and infected with *Botrytis cinerea*. *Eur. J. Plant Pathol.*, **106**: 657–665.
- Md. Shafiqur Rahman and M. N. Anwar. 2006. Fungitoxic and cytotoxic activity of a novel compound 1,2-benzenedicarboxylic acid, diisooctyl ester of *Plumbago zeylanica* linn. *Asian J of Micro., Biotech., and Environ. Sci.*, **8(3)**: 461-464
- Mehta, R., Radhakrishnan, T., Kumar, A., Yadav, R., Dobaria, J. R., Thirumalaisamy, P. P., Jain, R. K, and P. Chigurupati. 2013. Coat protein-mediated transgenic resistance of peanut (*Arachis hypogaea* L.) to peanut stem necrosis disease through *Agrobacterium*-mediated genetic transformation. *Indian J. Virol.* **24**: 205-213.
- Miao, Y., Zhu, L. and X. Zhang, 2019. Down regulation of cotton *GbTRP1* leads to accumulation of anthranilates and confers resistance to *Verticillium dahliae*. *J of Cotton Res.*, **2(1)**, 1-12.
- Mohammadi, M. and H. Kazemi. 2002. Changes in peroxidase and polyphenol oxidase activities in susceptible and resistant wheat heads inoculated with *Fusarium graminearum* and induced resistance. *Plant Sci.*, **162**, 491–498.
- Mohy El-Din, S. M. and M. M. Mohyeldin. 2018. Component analysis and antifungal activity of the compounds extracted from four brown seaweeds with different solvents at different seasons. *J of Ocean Uni of China.*, **17(5)**, 1178-1188.
- Moldovan, Mirela Carpa, Rahela Fizesan, Ionel, Vlase, Laurian Bogdan, Catalina Sonia, Iurian Benedec, Daniela Pop, Anca. 2020. Phytochemical Profile and Biological Activities of Tendrils and Leaves Extracts from a Variety of *Vitis vinifera* L. **9**. 1-20. 10.3390/antiox9050373.
- Morcia, C., Malnati, M. and V. Terzi. 2019. In vitro antifungal activity of terpinen-4-ol, eugenol, carvone, 1,8-cineole (eucalyptol) and thymol against mycotoxigenic plant pathogens. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess.*, **29(3)**:415-22.
- Morpurgo, R., Lopato, S.V., Afza, R. and F. J. Novak. 1994. Selection parameters for resistance to *Fusarium oxysporum* f.sp. *cubense* race 1 and race 4 on diploid banana (*Musa acuminata* Colla). *Euphytica* **75**, 121–129.
- Mozzetti, C., Farraris, L., Tamietti, G. and A. Matta. 1995. Variation in enzyme activities in leaves and cell suspension as markers of incompatibility in different *Phytophthora* pepper interaction. *Physiol. Mol. Plant Pathol.* **46**, 95–107.
- Munawwar, M. A., Chatta, F. A. and S. Kausar. 2018. Plant growth regulating activity of coumarin derivatives. *Horticult Int J.*, **2(2)**: 25-30.
- Pérez-Vicente, L., M. Dita and E. Martínez. 2014. Inoculation of *Fusarium oxysporum* f. sp. *cubense* causal agent of fusarium wilt in banana. Technical manual prevention and diagnostic of Fusarium wilt (Panama disease) of banana caused by *Fusarium oxysporum* f. sp. *cubense*. Tropical Race. 4: 55–58.

- Pintac, D., Majki, c, T., Torovi C, L., Beara, I. and N. Simin, 2018. Solvent selection for efficient extraction of bioactive compounds from grape pomace. *Ind. Crops. Prod.* **111**, 379–390.
- Ploetz, R. C. 2015. Management of Fusarium wilt of banana: a review with special reference to tropical race 4. *Crop. Prot.*; **73**: 7–15.
- Qi, D., Zou, L., Zhou, D., Chen, Y., Gao, Z., Feng, R. and W. Wang. 2019. Taxonomy and broad-spectrum antifungal activity of Streptomyces sp. SCA3-4 isolated from rhizosphere soil of *Opuntia stricta*. *Front in Micro.*, **10**: 1390.
- Razzaq, A., Sadia, B., Raza, A., Khalid Hameed, M. and F. Saleem. 2019. Metabolomics: A Way Forward for Crop Improvement. *J Metabolites*. **14;9(12)**:303.
- Robert, E and K. Helmut. 1992. Isoflavonoid phytoalexins and their biosynthetic enzymes. In: Sarah, J.G., McPherson, D.J., Bowles, D.J. (Eds.), *Mol Plant Pathology*, vol. **11**. IRL, Oxford, England, pp. 45–62
- Roberts, M., Minott, D. A., Tennant, P. F. and J. C. Jackson. 2008. Assessment of compositional changes during ripening of transgenic papaya modified for protection against papaya ringspot virus. *J of the Sci of Food and Agri.*, **88(11)**: 1911-1920.
- Shomo, S. H. Shifa, C. Senthilraja, Madem Gurivi Reddy, V. Paranidharan and R. Velazhahan. 2016. Analysis of metabolite profiles in leaf tissues of transgenic groundnut plants modified for resistance against Tobacco streak virus. *Biochem. Cell. Arch.*, **16: (2)**: 245-250.
- Shobier, A.H.S.; Abdel Ghani, S.A. and K. M. Barakat. 2016. GC/MS spectroscopic approach and antifungal potential of bioactive extracts produced by marine macroalgae. *Egypt J. Aquat. Res.*, **42**: 289–299
- Subramaniam, S., Maziah, M., Sariah, M., Puad, M. P. and R. Xavier. 2006. Bioassay method for testing Fusarium wilt disease tolerance in transgenic banana. *Sci Horticulturae*, **108(4)**: 378-389.
- Sunisha, C., Sowmya, H. D., Usharani, T. R., Umeha, M., Gopalkrishna, H. R. and A. Saxena. 2020. Deployment of stacked antimicrobial genes in banana for stable tolerance against *Fusarium oxysporum* f. sp. *cubense* through genetic transformation. *J Molecular biotechnology*, **62(1)**: 8-17.
- Waite, B. H. and R. H. Stover, 1960. Studies on Fusarium wilt of bananas. VI. Variability and the cultivar concept in *Fusarium oxysporum* f. *cubense*. *J. Botany*; **38**, 985-994.
- Walters, D. R., Walker, R. L., and K. C. Walker. 2003. Lauric acid exhibits antifungal activity against plant pathogenic fungi. *J of Phytopathology*, **151(4)**: 228-230.
- Yang, Y., Chen, Y., Cai, J., Liu, X. and G. Huang. 2021. Antifungal activity of volatile compounds generated by endophytic fungi *Sarocladium brachiariae* HND5 against *Fusarium oxysporum* f. sp. *cubense*. *J PLOS one*, **16(12)**: e0260747.
- Yuan, J., Raza, W., Shen, Q. and Q. Huang. 2012. Antifungal activity of *Bacillus amyloliquefaciens* NJN-6 volatile compounds against *Fusarium oxysporum* f. sp. *cubense*. *Applied and Environmental Micro.*, **78(16)**: 5942-5944.
- Zishan, M. and U. Manzoor. 2020. Antifungal Activity of *Aloe Barbadensis* and *Nigella Sativa*: A Review. *Int J of Biological, Physical and Chemical Studies*, **2(2)**, pp.11-14.