



RESEARCH ARTICLE

Response of Sesamum Varieties to Graded Levels of Sulphur on Seed Yield and Oil Content in Soils of Different Sulphur Status

Venkatakrishnan L^{1*}, Backiyavathy M R², Meena S¹, Kalaiselvi T² and Amirtham D³

¹Department of Soil Science & Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore - 641 003.

²Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore - 641 003.

³Department of Food Processing and Post-Harvest Technology, Tamil Nadu Agricultural University, Coimbatore - 641 003.

ABSTRACT

Field experiments were conducted in Coimbatore District with discrete soil sulphur status i.e., soils with sufficient and deficient levels of sulphur, to know the relative response of graded levels varied doses of sulphur on two different sesamum varieties. The experiments were laid out in a split-plot design with two factors viz., two sesamum varieties (TMV 7 and SVPR 1) and six sulphur levels (0, 20, 30, 40, 50, and 60 kg S ha⁻¹). The results revealed that application of 30 and 50 kg S ha⁻¹ at sulphur sufficient and sulphur deficient soils, respectively, has increased the yield parameters such as test weight, seed weight per plant, seed yield, stalk yield, oil yield, and biological yield. Between two varieties, TMV 7 has performed better when compared with SVPR 1. Under both the soil sulphur status, further increase in the level of sulphur recorded a nonsignificant increase in the yield attributes.

Received : 28th October, 2020

Revised : 07th November, 2020

Revised : 14th November, 2020

Accepted : 28th November, 2020

Keywords: *Sulphur levels; Sulphur status; Growth; Yield attributes*

INTRODUCTION

Sesame is commonly known as “Till.” Its botanical name is *Sesamum indicum* L. It is one of the earliest domesticated plants. Due to the presence of a potent antioxidant, sesame seeds are known as “the seed of immortality”. Two distinct types of seed are recognized, the white and the black. There are also intermediate colored varieties varying from red to rose or from brown or grey.

India is the foremost producer of oilseed crops in terms of area (26 million ha) and production (30 million tonnes), in which 72% of the area is under rainfed condition. As per 4th Advance Estimates Oilseeds Production in the country during 2017-18, the production was estimated as 31.31 million tonnes which is marginally higher than the production of 31.28 million tonnes during 2016-17. However, the production of oilseeds during 2017-18 was higher by 1.76 million tonnes than the average of 2012-13 to 2016-17. The per capita consumption was 15.80 kg per person per annum during 2012-13 and has increased to 19.30 kg per person in 2017-18. In addition, demand for edible oils in both quantity and quality is growing due to population growth, which was 1.21 billion in 2011 (Luna, 2011) and now reached above 1.25 billion (Gupta, 2014). Considering the growing domestic demand for edible oils, it has now been planned to achieve a production of 45.64 million tonnes from nine annual

oilseed crops by 2022-2023, expecting an additional production of about 15.58 million tonnes over and above the 30.06 million tonnes production during quinquennium ending 2016-17. Thus, the availability of total vegetable oil from domestic production would be about 13.69 million tonnes by 2022 (30 % recovery) as against the current annual output of about 7.0 million tonnes (NFSM, 2017).

Among the oilseed, sesame is one of the most important oil seed crops belonging to the Pedaliaceae family and extensively grown in different parts of the world and it ranks fourth among oil seed crops of the world in terms of area of production. It is a versatile crop with a diversified use of high-quality edible oils that contains vitamins, amino acids and polyunsaturated fatty acids. In sesame seeds, linoleic and oleic acids are the prime factors, which are responsible for oil consistency (Uzun *et al.*, 2008).

Generally, the sulphur (S) requirement by crops is equal to that of phosphorus, which plays a vital role in plant metabolism. It is indispensable for the synthesis of essential oils (Singh *et al.*, 2000), a constituent of several organic compounds (Shamina and Imamul, 2013), oil storage organs, particularly oil glands i.e., glyoxysomes (Jaggi *et al.*, 2000) and vitamin B1 (Thirumalaisamy *et al.*, 2001). It

*Corresponding author's e-mail: krishnann93@gmail.com

also serves as a component of plant amino acids, proteins, vitamins, and enzyme structures. It has been observed that increasing sulphur application increases oil content and oil yield, protein, and glucosinolates of canola seeds (Haneklaus *et al.*, 1999). Sulphur also influences the productivity of oil seed and total oil content (Egesel *et al.*, 2009). In addition, the continuous application of NPK fertilizers contributes to a sulfur deficiency along with a great deal of organic matter deficiency, making the situation worse for oilseed crops. In view of this background, the present study was taken up to trace the optimum level of sulfur recommendation for yield maximization and quality improvement in sesamum.

MATERIAL AND METHODS

Two field experiments were conducted at two different locations of Coimbatore district in soils having two sulphur status i.e., a soil with sufficient level and soil with deficient level of sulphur in order to study the response of two sesamum varieties to graded doses of sulphur with respect to seed yield and quality. The field experiments were conducted at farmers' holdings in Selambanur (Latitude: 11.00° N and Longitude: 76.79E) and Puthur (Latitude: 10.979436 and Longitude: 76.84E) villages of Thondamuthur block at Coimbatore district from February 2019 to July 2019 and March 2019 to August 2019, respectively. The selected experimental field soils were neutral in pH 7.5 & 7.2 and has an electrical conductivity of 0.19 & 0.15 dSm⁻¹, available N of 361 & 349 kg ha⁻¹; available P of 19 & 21 kg ha⁻¹; available K of 295 & 288 kg ha⁻¹; available S of 43.23 & 7.3 mg kg⁻¹ respectively

in soils of high and low sulphur status field. The treatments comprised six levels of sulphur supplied using gypsum as the source (0, 20, 30, 40, 50 and 60 kg S ha⁻¹) along with soil test based application of major nutrients fertilizer doses, which were replicated three times in a split-plot design (Main plot – sesamum varieties, Sub plot- S levels). The data on yield parameters, seed yield and oil content were analyzed statistically using the AgRes statistical software (Pascal Intel Software Solutions). Wherever the treatment differences were found significant, critical differences (CD) were worked out at 5% level of significance with mean separation by least significant difference and denoted by the symbol (* for 5% and ** for 1%). A simple correlation was worked out between different parameters to know the relationship that exists among them (Gomez and Gomez, 1984).

RESULTS AND DISCUSSION

Test weight

Statistically significant variations were recorded for sulphur application with respect to the test weight of sesame seeds. The test weight of sesamum varieties with 30 kg S ha⁻¹ was significantly higher in the case of sulphur sufficient soil. In the case of sulphur deficient soil, the highest test weight of 3.51 g (TMV 7) and 3.45 g (SVPR 1) were recorded by applying 50 kg S ha⁻¹ which was significantly higher than all the other treatments. Among the varieties, TMV 7 has recorded the highest test weight and proved to be the best responding variety than SVPR 1 (Table 1).

Table 1. Effect of graded levels of sulphur on test weight of sesamum varieties

Treatments	Test weight (g)					
	Sulphur sufficient soil			Sulphur deficient soil		
	TMV 7	SVPR 1	Mean	TMV 7	SVPR 1	Mean
Absolute Control	2.41	2.29	2.35	2.10	2.00	2.05
RDF alone	2.62	2.51	2.57	2.31	2.21	2.26
RDF+ S 20 kg ha ⁻¹	2.84	2.72	2.78	2.53	2.45	2.49
RDF+ S 30 kg ha ⁻¹	3.11	3.02	3.07	2.75	2.67	2.71
RDF+ S 40 kg ha ⁻¹	3.14	3.05	3.10	2.97	2.89	2.93
RDF+ S 50 kg ha ⁻¹	3.17	3.10	3.14	3.51	3.45	3.48
RDF+ S 60 kg ha ⁻¹	3.15	3.11	3.13	3.28	3.25	3.27
Mean	2.92	2.83		2.78	2.70	
	SEd	CD		SEd	CD	
V	0.006	0.02**		0.01	0.05*	
T	0.035	0.07**		0.05	0.10**	
V at T	0.047	NS		0.06	NS	
T at V	0.050	NS		0.07	NS	

*RDF –Recommended dose of NPK fertilizers

The increase in test weight might be due to the role of sulphur in activating the growth and yield components (Tahir *et al.*, 2014; Ojoni *et al.*, 2018).

Seed yield

The effect due to varieties, sulphur levels and

their interactions were prominent in case of seed yield (Table 2). In sulphur sufficient soil, the seed yield of varieties were significantly higher by applying 30 kg S ha⁻¹ which was on par with 40, and 50 kg S ha⁻¹, however application of 60 kg S ha⁻¹ showed decrease in seed yield.

Table 2. Effect of graded levels of sulphur on seed yield of sesamum varieties

Treatments	Seed yield (kg ha ⁻¹)					
	Sulphur sufficient soil			Sulphur deficient soil		
	TMV 7	SVPR 1	Mean	TMV 7	SVPR 1	Mean
Absolute Control	652	573	613	619	519	569
RDF alone	725	633	679	643	542	593
RDF+ S 20 kg ha ⁻¹	801	696	749	672	574	623
RDF+ S 30 kg ha ⁻¹	870	760	815	733	638	686
RDF+ S 40 kg ha ⁻¹	876	766	821	794	692	743
RDF+ S 50 kg ha ⁻¹	880	770	825	846	754	800
RDF+ S 60 kg ha ⁻¹	860	765	813	836	736	786
Mean	809	709		735	636	
	SEd	CD		SEd	CD	
V	7.7	33.2**		5.5	23.9**	
T	13.1	27.1**		12.1	25.0**	
V at T	18.8	NS		16.8	NS	
T at V	18.6	NS		17.1	NS	

Whereas in case of sulphur deficient soil, the highest seed yield was recorded with the application of 50 kg S ha⁻¹ irrespective of the varieties. Considering the varietal effects, TMV 7 has proven to be the best responding variety than SVPR 1. The increase in seed yield could be attributed to the promotion of photosynthesis, net assimilation rate and crop growth rate due to the S application, which resulted in higher seed yield (Tahir *et al.*, 2014; Suchhanda *et al.*, 2016).

application at the rate of 30 kg S ha⁻¹. Further increase of sulphur application did not increase the yield significantly. However, in sulphur deficient soil, a significant increase in stalk yield was recorded up to 50 kg S ha⁻¹ with a yield of 3175 and 3154 kg ha⁻¹. Among the varieties, TMV 7 has recorded the highest stalk yield than that of SVPR 1. Suchhanda *et al.* (2016) and Ojoni *et al.*, (2018) have reported a similar increase in stalk yield.




Figure 1. Effect of graded levels of sulphur application on oil content

Stover yield

The stalk yield of sesamum was significantly influenced by graded levels of sulphur application (Table 3). In sulphur sufficient soil, the stalk yield has shown a significant increase with sulphur




Figure 2. Effect of graded levels of sulphur application on biological yield

Oil content

The graded levels of sulphur application (Figure 1) significantly influenced the oil content of different varieties. The oil content has shown a significant increase by applying sulphur up to 30 kg S ha⁻¹; the increase in oil content was not significant with

a further increase of sulphur application in sulphur sufficient soil. However, in the case of sulphur deficient soil, a significant increase in oil content was recorded up to 50 kg S ha⁻¹ with a yield of 49.90 per cent and 52.30 per cent, respectively, for TMV 7 and SVPR 1. Among the varieties, SVPR 1 has

shown increased oil content. This might be due to the vital role of sulphur in the synthesis of S-containing amino acids and vitamins like biotin, thiamine, vitamin B1 as well as the formation of ferredoxin, an iron-containing plant protein that is required for the production oil. Bosale *et al.* (2011) and Dharati *et al.* (2017) also obtained similar results.

Table 3. Effect of graded levels of sulphur on stover yield of sesamum varieties

Treatments	Stover yield (kg ha ⁻¹)					
	Sulphur sufficient soil			Sulphur deficient soil		
	TMV 7	SVPR 1	Mean	TMV 7	SVPR 1	Mean
Absolute Control	3034	2975	3005	2960	2951	2956
RDF alone	3067	3025	3046	2994	2983	2989
RDF+ S 20 kg ha ⁻¹	3110	3086	3098	3024	3013	3019
RDF+ S 30 kg ha ⁻¹	3195	3175	3185	3065	3052	3059
RDF+ S 40 kg ha ⁻¹	3205	3184	3194	3085	3061	3073
RDF+ S 50 kg ha ⁻¹	3211	3193	3202	3175	3154	3165
RDF+ S 60 kg ha ⁻¹	3204	3184	3194	3165	3163	3164
Mean	3146	3117		3067	3054	
	SEd	CD		SEd	CD	
V	12.8	NS		8.07	NS	
T	39.0	80.5**		38.4	79.2**	
V at T	52.6	NS		50.9	NS	
T at V	55.1	NS		54.3	NS	

Biological yield

Data presented in Figure 2 showed that application of sulphur to sesamum had significantly enhanced the biological yield at different soil sulphur status. In the case of sulphur sufficient soil, the yield response was significantly increased up to 30 kg S ha⁻¹ and on further increase in sulphur level through gypsum has resulted in a non-significant trend. In accordance to the sulphur application, in sulphur deficient soil, the response of yield was extended up to 50 kg S ha⁻¹ with a yield of 4021 and 3908 kg ha⁻¹ respectively for two varieties, among which TMV 7 had produced a higher biological yield than SVPR 1.

CONCLUSION

Based on the experiment conducted in soils of two different sulphur statuses with two different sesamum varieties, it is concluded that TMV 7 proves to be a better responding variety than that of SVPR 1 by applying sulphur @ 30 and 50 kg ha⁻¹ in sulphur sufficient and deficient soils. The results indicated that supplementation of sulphur nutrition to sesamum in sulphur deficient soil is a boon for the increased growth and yield of sesamum crop.

REFERENCES

Bhosale, N.D., Dabhi, B.M., Gaikwad V.P. and A.B. Chavan. 2011. Growth, yield and quality parameters of sesamum (*Sesamum indicum* L.) as influenced

- by different levels of potash and sulphur. *Inter. J. Forestry & Crop Improv.*, **2(20)**: 121-123.
- Dharati P Patel., Patel, R.A., Sonaka Ghosh and P.K.Patel. 2017. Effect of irrigation, Vermicompost and sulphur on oil content and S content and uptake of summer sesamum (*Sesamum indicum* L.) under loamy sand. *Inter. J. of Chem. Studies.*, **5(5)**: 2159-2162
- Egesel., Kemal Gül M. and FatihKahr Aman. 2009. Changes in yield and seed quality traits in rapeseed genotypes by sulphur fertilization. *Eur Food Res Technol.* **229**:841
- Gomez Kwanchai, A. and Arturo Gomez. 1984. Statistical Procedures for Agricultural Research: John Wiley & Sons.
- Gupta MP. 2014. Challenges of food security. *The Daily Excelsior.*, **50(77)**:6.
- Haneklaus S, Paulsen, H.M., Gupta, A.K., Bloem, E. and E.Schnug. 1999. Influence of sulphur fertilization on yield and quality of oilseed and mustard. In: Proc. 10th Int. Rapeseed Cong., Canberra, Australia.
- Jaggi, R., Rakesh, C., Sharma, K. and S.S. Paliyal. 2000. Sulphur as second major nutrient in cruciferous oilseeds. *Indian Farming.* **12**:14-18.
- Luna RK. 2001. How India can reap demographic dividend. *The Tribune.*, **131(103)**:11.
- NFSM., 2017. Present Status of Oilseed crops and vegetable oils in India, Status paper on oil seeds Ministry of Agriculture and Farmers welfare. <https://www.nfsm.gov.in/StatusPaper/NMOOP2018.pdf>.

- Ojoni Tekseng., Sing,P.K. and Kevineituo Bier. 2018. Effect of phosphorus and sulphur nutrition on growth, yield and quality attributes of sesame (*Sesamum indicum* L.) under acidic soil of Nagaland. 315-319.
- Scherer, H.W. 2001. Sulphur in crop production. *Eur. J. Agron.*, **14**:81-111.
- Shamina., Imamul. 2003. Mineralization pattern of added sulphur in some Bangladesh soils under submerged condition. *Indian J. Agric. Chem.*, **36**: 13-21.
- Singh A., Singh, S.P., Katiyar, R.S. and P.P. Singh. 2000. Response of nitrogen and sulphur on economic yield of sunflower under sodic soil condition. *Indian J. Agric. Sci.*, **70**: 536-537.
- Suchhanda Mondal. 2016. Efficiency of sulphur source on sesame (*Sesamum indicum* L.) in red and lateritic soil of West Bengal. *Inter. J. of plant, animal and environmental sciences.*, **6(2)**: 65-70.
- Tahir, M., Ibrahim,A., Tahir, S., Aimen Ayub ., Tanveer,A. and H. Rehman. 2014. Effect of sulfur levels on two sesame (*Sesamum indicum* L.) varieties under climatic conditions of pakistan. *Inter. J. of plant & soil science.*, **3(3)** : 281-288.
- Thirumalaisamy, K., Srinivasan, P.S., Pangarusamy, U., Thangaraj, M., Mallika, V. and S. Kannaiyan . 2001. Nutrient deficiency and remedial management in crops. 1st Edn.,Sigar Graphics Pub. Coimbatore, London. 10-11.
- Uzun, B., Arslan, C. and S. Furat . 2008. Variation in fatty acid compositions, oil content and oil yield in a germplasm collection of sesame (*Sesamum indicum* L.). *J Am Oil Chem Soc.*, **85**: 1135–1142.