



RESEARCH ARTICLE

## RNAi Mediated Silencing of *ftf1* Gene Enhances Resistance in Banana to Fusarium Wilt

Deepa Sankari R<sup>1</sup>, Kumar K K<sup>2</sup>, Varanavasiappan S<sup>2</sup>, Kalpana K<sup>1</sup>, Sunisha C<sup>6</sup>, Usharani T R<sup>6</sup>, Muthulakshmi P<sup>4</sup>, Auxcilia J<sup>5</sup>, Arul L<sup>2</sup>, Kokiladevi E<sup>2</sup>, Nakkeeran S<sup>3</sup>, Sudhakar D<sup>2\*</sup> and Eraivan Arutkani Aiyanathan K<sup>1\*</sup>

<sup>1</sup>Department of Plant Pathology, Agricultural College and Research Institute, Madurai - 625 104

<sup>2</sup>Department of Plant Biotechnology, Centre for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University, Coimbatore-641003

<sup>3</sup>Agricultural College and Research Institute, Kudumiyamalai-622 104

<sup>4</sup>Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore-641003

<sup>5</sup>Department of Fruit Science, Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore-641003

<sup>6</sup>Division of Biotechnology, ICAR- Indian Institute of Horticultural Research, Hesarghatta Lake post, Bangalore - 560 089

### ABSTRACT

Fusarium wilt caused by *Fusarium oxysporum f. sp. cubense* (Foc) is a devastating disease of banana. Race 1 of Fusarium wilt is a serious threat to *Musa* spp. cv. Rasthali (AAB, Silk subgroup), which is a choice variety traditionally grown in most of the South East Asian countries. The investigation was an attempt to study whether RNAi-mediated expression of small interfering RNAs (siRNAs) targeted against *ftf1* (fusarium transcription factor1) gene in transgenic banana could impart resistance against Foc race 1. Partial sequences comprising sense and antisense sequences of *ftf1* gene were assembled as siRNA construct in a binary vector background and transformed into embryogenic cell suspensions of banana cv. Rasthali by *Agrobacterium* mediated genetic transformation. Presence of RNAi construct in transgenic banana plants were confirmed by polymerase chain reaction assay. Forty- six transformed lines derived from RNAi-*FTF1* were found to be free of external symptoms of Foc after 3 months of challenge inoculation. Seventeen transgenic lines continued to resist the Foc race 1 even after six months of post inoculation. The present study confirmed that the host induced RNAi-mediated gene silencing of a fungal gene could confer resistance against Foc in banana.

Received: 22 June 2022

Revised: 09 August 2022

Accepted: 08 September 2022

**Keywords:** *Banana Fusarium wilt; Fusarium oxysporum f. sp. cubense; RNA interference; fusarium transcription factor 1*

### INTRODUCTION

Banana (*Musa spp.*) is one of the most important fruit crops as well as staple food crop for millions of people in developing countries. Most of the edible bananas are triploid, which are formed by inter-specific hybridization between two diploid progenitors, namely *Musa acuminata* (AA genome) and *Musa balbisiana* (BB genome). A major threat to banana production is Panama disease or Fusarium wilt caused by a soil borne pathogen, *Fusarium oxysporum f.sp. cubense* (Foc). Foc invades banana root system, colonizes the corm tissue and clogs the water conducting vessels, thereby causing wilting of the aerial parts of infected banana plant. The infected plant shows typical discoloration of the corm and pseudostem and yellowing of the foliage followed by complete wilting (Dale et al., 2017).

The Fusarium wilt was first discovered in banana var. Sugar (Silk AAB) in Brisbane, Queensland, Australia during 1876 (Bancroft, 1876). The pathogen has been classified into four races, namely races 1, 2, 3 and 4. Race 1 which infected Gros Michel (AAA) earlier, now infects Silk and Pome (AAB) and Pisang Awak (ABB); race 2 infects cooking bananas namely Bluggoe (ABB) subgroup; race 3 infects ornamental namely *Heliconia sp.*, (Tropical American banana relatives) and race 4 affects all cultivars in the Cavendish (AAA) subgroup in addition to the varieties susceptible to races 1 and 2 (Buddenhagen, 2009; Chittarath et al., 2018; Zheng et al., 2018; Thangavelu et al., 2021). Rasthali cultivar of banana (AAB) is exceptionally valued for its unique fragrance, taste and keeping quality. Moreover, it is a premium variety cultivated across the globe, more extensively in the South East Asian countries. Unfortunately, Rasthali is highly susceptible to Fusarium wilt (race 1) resulting in heavy yield loss to the farmers. In India, besides

\*Corresponding author's e-mail: [deepasankariagri02@gmail.com](mailto:deepasankariagri02@gmail.com)

Rasthali, the major varieties affected are Amrithapani, Karpuravalli, Monthan, Ney Poovan, and Virupakshi (Mustafa and Thangavelu, 2010).

Various agronomic practices like crop rotations, addition of organic amendments, flooding the field etc., have proved to be ineffective in controlling this disease (Bakry *et al.*, 2009). Once the fungal spores get established in field, they persist in soil for almost three decades. Moreover, the disease is difficult to manage using fungicides (Thangavelu and Mustafa, 2010). One of the sustainable solutions to manage this disease is to develop resistant cultivars by either conventional breeding or genetic engineering. Conventional breeding in banana faces challenges due to various sexual reproduction barriers such as high sterility, complex genetic background, polyploid and parthenogenesis (Ghag *et al.*, 2014), making it extremely difficult to develop new disease resistant bananas (Czislowski *et al.*, 2018). Therefore, host-induced gene silencing (HIGS) has increasingly become one of the most practical approaches for enhancing new disease resistant crop varieties (Qi *et al.*, 2019; Zhang *et al.*, 2016).

RNA interference (RNAi), also known as Host-Delivered RNAi (HD-RNAi) or Host Induced Gene Silencing (HIGS) or pathogen derived resistance, uses the host plant as a delivery system to cause gene silencing in the pathogen (Nunes and Dean, 2012; Ghag *et al.*, 2014; Mat Jalaluddin *et al.*, 2019). Plant siRNAs are 21 to 24 nucleotide long double-stranded RNAs with 3' 2-nucleotide overhangs and generally carry a 2'-O methyl group on the 3'-terminal ribose (Yu *et al.*, 2010). Double stranded RNAs, the progenitors of siRNAs are processed by dicer proteins in the nucleus and then the siRNAs are incorporated into the RNA-induced silencing complex (RISC) in the cytoplasm and thus results in sequence specific mRNA degradation (Bernstein *et al.*, 2001; Baulcombe, 2004; Ghildiyal and Zamore, 2009). Hemi biotrophic fungus like Foc have a short biotrophic phase followed by all out necrotrophy of the targeted plant. During the short biotrophic phase, the fungal haustoria penetrate the cell wall to derive nutrition, activating plant necrosis through a cascade of plant signaling system (Thaler *et al.*, 2004). It is probably during this short biotrophic phase, exchange of cellular components occurs between host and the pathogen. The host derived siRNAs (targeted against vital fungal genes) finds entry into the fungal cytosol leading to host-induced post-transcriptional gene silencing of the target fungal genes.

Genetic engineering employing RNA interference technology has been successfully used to incorporate resistance against viruses (Shekhawat *et al.*, 2012), nematodes (Huang *et al.*, 2006), insect pests (Mao *et al.*, 2007) and fungal

pathogens (Nowara *et al.*, 2010; Panwar *et al.*, 2013; Ghag *et al.*, 2014 and Sunisha *et al.*, 2020). The *ftf1* gene encoding a Zn (II) 2-Cys6 binuclear DNA-binding protein is up-regulated several folds in *Fusarium oxysporum f. sp. phaseoli* during its colonization in common bean (De Vega-Bartol *et al.*, 2011). This indicated that the down regulation of *ftf1* gene could potentially impart resistance to Fusarium. Besides *ftf1* gene, velvet gene was reported to play a key role in virulence of the pathogen. Down regulation of *ftf1* or velvet gene could confer resistance to Fusarium in banana (Ghag *et al.*, 2014). Considering these leads, an attempt was made to develop Foc race 1 resistant transgenic banana in Rasthali by silencing *ftf1* (fusarium transcription factor 1) gene.

## MATERIAL AND METHODS

### Construction of RNAi vector

Partial sequence of *ftf1* gene (323 bp) was amplified from genomic DNA of Foc race 1 as described by Ghag *et al.* (2014) using gene specific primers designed to create a restriction site on either side of the amplified product (Table 1). The amplified products were cloned between XhoI and XbaI sites of pJET vector. The cloned sequences were confirmed by sequencing. The partial *ftf1* sequence (antisense sequence) was cloned between KpnI and SpeI (New England Biolabs, UK) sites of an intermediate vector, pSTARLINGA followed by cloning of sense sequence between BamHI & AscI sites. Finally, the RNAi cassette (*ftf1*sense-intron-*ftf1*antisense) (Figure. 1) was released from recombinant pSTARLINGA using NotI and subcloned in the NotI site of a binary vector (pART27). The vector was then mobilized into *Agrobacterium tumefaciens* strain, AGL1 by triparental mating. *Agrobacterium* strain, AGL1 harboring recombinant pART27 was used for genetic transformation of Rasthali.




Figure 1. Physical map of pART27 harboring *ftf1* RNAi cassette

### Generation and molecular analysis of transgenic banana plants

A single colony of AGL1 strain harboring *ftf1* gene was inoculated in YEMA (Yeast Extract Maltose Agar) liquid medium added with 100 mg L<sup>-1</sup> spectinomycin and grown in an orbital shaker for 12 h at 28 °C. An overnight grown *Agrobacterium* culture (1 mL) was transferred to 50 mL of AB minimal medium augmented with 100 mg L<sup>-1</sup> spectinomycin and incubated under the same conditions. The *Agrobacterium* culture was pelleted (3500 rpm for 10 min), suspended in AB induction medium with 100 mM acetosyringone (ACS), and allowed to grow at 28 °C (90 rpm for 3-4 h). Following induction, the bacterial culture was centrifuged (3500 rpm for 10 min) and resuspended in the same medium with the addition of pluronic acid F68 (0.02 %) to a final OD600 nm of 0.5.

**Table 1. List of primers used to amplify target genes**

Name of gene	Forward (F) and reverse (R) primers (5' to 3')	Amplicon size (bp)	PCR conditions
Partial <i>ftf1</i> sense and antisense	Antisense <i>ftf1</i> F* CTTCGGTACCCCTTG AAGCTAGTCTGCCC TTG Antisense <i>ftf1</i> R* GAAGACTAGTGAG GAAAAGGCGCAGT ATCCA Sense <i>ftf1</i> F** TGTTGGATCCCTTG AAGCTAGTCTGCCC TTG Sense <i>ftf1</i> R** AACAGGCGCGCGG AGGAAAAGGCGCA GTATCCA	323	94 °C for 5 min: 1 cycle 94 °C for 1 min 54 °C for 30sec 72 °C for 30 sec 72 °C for 10 min: 1 cycle 35 cycles
<i>ftf1</i>	<i>ftf1</i> F AGTCTGCCTTGTG ACTCAA <i>ftf1</i> R AGGAAAAGGCGCAG TATCCAG	287	94 °C for 5 min: 1 cycle 94 °C for 1 min 54 °C for 30sec 72 °C for 30 sec 72 °C for 10 min: 1 cycle 35 cycles
<i>nptII</i>	<i>nptII</i> F CTGATGCTCTTCGTCCAG AT <i>nptII</i> R AGAGGCTATTCGGCT ATGACT	440	94 °C for 5 min: 1 cycle 94 °C for 1 min 63 °C for 40sec 72 °C for 45 sec 72 °C for 10 min: 1 cycle 30 cycles
SIX6B	SIX6B F ACGCTTCCCAATACCGTC TGT SIX6B R AAGTTGGTGAGTATCAAT GC	220	94 °C for 5 min: 1 cycle 94 °C for 1 min 63 °C for 40sec 72 °C for 45 sec 72 °C for 10 min: 1 cycle 30 cycles

The transformation was performed using the protocol as suggested in technical guideline bulletin 8 of INIBAP (Stross *et al.*, 2003) with few modifications. About 0.5 ml of settled cell volume (SCV) of banana embryogenic cells were subjected for heat shock at 45 °C for 5 min followed by co-cultivation with *Agrobacterium* by centrifugal method. The co-cultivated cells were inoculated on co-cultivation medium (CCM) with 100 µM ACS and incubated for 3 days at 28 °C. The cells were

washed in MA2 liquid medium and plated on semi-solid MA2 medium supplemented with cefotaxime (250 mg L<sup>-1</sup>) for 2 weeks. Cells were transformed on MA3 medium (250 mg L<sup>-1</sup> of cefotaxime and 100 mg L<sup>-1</sup> of kanamycin) for maturation of somatic embryos. Matured embryos were then plated on MA4 medium supplemented with BAP 0.5 mg L<sup>-1</sup> for shoot regeneration. The rooting was achieved on MA5 medium supplemented with 1 mg L<sup>-1</sup> BAP and 2 mg L<sup>-1</sup> IBA. For plantlet acclimatization, the rooted plants were hardened in greenhouse and maintained further for molecular analysis.

All the transgenic lines were further confirmed using primers specific for the neomycin phosphotransferase II gene and *ftf1* gene sequences (Table 1). Genomic DNA was isolated from leaves of transgenic and non-transformed banana plants using CTAB method (Dellaporta, 1983). The conditions for PCR were as follows: initial denaturation at 94 °C for 4 min; denaturation at 94 °C for 1 min; annealing at 57 °C for 45 s; extension at 72 °C for 1 min and 35 cycles were performed before the final extension at 72 °C for 10 min.

### Screening of transformants for resistance to *Foc* race 1

A highly virulent fungal strain (*Foc* race 1), KP used in this study was maintained in PDA plates. For inoculum production, 500 mL Erlenmeyer flasks containing 250 mL of potato dextrose liquid medium were inoculated with 9 mm dia mycelial plugs of *Foc* race 1 excised from 5-day-old colonies grown on Potato Dextrose Agar (PDA) medium and incubated for 6 days (28±2 °C, 200 rpm). In addition, *Foc* mycelial discs were grown in 250 mL flasks containing 50 g of sterilized bajra kernels (28±2 °C, 10 days) as described by Dita *et al.* (2010).

Screening for resistance against *Fusarium* wilt of banana was conducted in transgenic events under contained conditions, using non-transgenic Rasthali control as susceptible and Grand naine as resistant cultivar. To evaluate fungal resistance, transgenic and control plants were inoculated with *Foc* race 1 strain, KP at five-leaf stage. Prior to inoculation, plants were removed from pots and the roots were trimmed to a length of ~40 cm. Inoculation was performed by root dipping (30 min, 106 conidia mL<sup>-1</sup>) and inoculated plants were transferred to pots, watered with tap water daily and supplemented with 50 ml of Hoagland's solution at weekly interval. The disease severity of each plant was scored at 45 days and 3 months post inoculation. For resistance evaluation, the standard evaluation method described by Perez-Vicente *et al.* (2014) was adopted. The following scale was used to phenotype the plants based on the percentage of yellowing of leaves: 0 = no symptoms; 1 = initial yellowing, mainly in the lower leaves; 2 = yellowing of all the lower leaves with some discoloration of younger leaves;

3 = all leaves with intense yellowing or the plant died. Both transformed and non-transformed plants were scored based on their symptoms.

### Detection of *Foc* in transgenic plants by PCR

Total genomic DNA was isolated from infected plants using the CTAB method as stated above. PCR analysis was done using *Foc* race 1 specific SIX6B gene primers to detect the presence of *Foc* (Table 1).


## Results and Discussion

The production and trade of banana is threatened by major fungal disease such as Fusarium wilt which can damage huge plantations in a short span of time. Banana cv. Rasthali (AAB) is considered to be highly susceptible to this disease and is threatened with extinction (Thangavelu *et al.*, 2001). In banana, a vegetatively propagated crop with triploid genome, recombination breeding has not been successful in introducing resistance from tolerant genomes. RNAi phenomenon is being exploited in plants for resistance against biotic stresses that affects crop productivity significantly and this technology has been recognized as key biotechnology strategy in plant stress management. The expression of dsRNA, targeting essential genes of pathogens to enhance resistance against a number of pathogens have been reported in several crop plants including wheat, barley, banana and tomato (Nowara *et al.*, 2010; Ghag *et al.*, 2012, 2014; Singh *et al.*, 2020). The regulatory approved Brazilian transgenic bean employing RNAi technology to combat bean golden mosaic virus is an example. Realizing the potential of the RNAi technology, this approach was used to engineer expression of dsRNA specific to Fusarium transcription factor 1 gene in banana plants with a view to inhibit the gene expression. The production of fungal gene-specific siRNAs in banana plants was expected to inhibit the invasion of host cells by the fungus and thereby reducing disease severity. An RNAi vector in the backbone of binary plasmid, pART27 was constructed and mobilized into *Agrobacterium tumefaciens* and used for genetic transformation of embryogenic cell suspension of Rasthali.


### Generation and molecular analysis of transgenic banana plants expressing RNAi constructs

Co-cultivation of embryogenic cell suspension cultures of banana cv. Rasthali using *Agrobacterium tumefaciens* AGL1 strain harboring pART27-RNAi-*FTF1* resulted in 444 plantlets (Figure 2a-e). Genomic DNA of all the putatively transformed plants and plants derived from non-transformed ECS were used as template in CR using primers specific to the neomycin phosphotransferase II and *ftf1* gene. Amplification of internal sequences of 440 bp of *nptII* and 287 bp of *ftf1* gene was observed in 441 plants

derived from RNAi-*FTF1* construct, whereas the same was absent in non-transformed controls (Figure 3 a & b).



**Figure 2. Transformation of banana cv. Rasthali.** Embryogenic cells were co-cultivated and plated on co-cultivation medium, b) developed somatic embryo cells on MA3 medium, c) germination of plantlets from the somatic embryos on MA4 medium, d) rooted plants and e) putative transformants acclimatized in protray



**Figure 3. PCR analysis of putative transgenic banana cv. Rasthali.** Lane 1- 1 kbp marker, lane 2- positive control (pART27 *ftf1*), lane 3-negative control (Rasthali plant DNA), lane 4-water control, lanes 5 to 20-banana transformants

### Bioassay of transgenic plants against *Foc* race 1

Transgenic banana plants (434 plants) harboring RNAi-*ftf1* construct were subjected to fungal bioassay following acclimatization for 2 months in greenhouse. Transgenic plants were found to be apparently healthy and exhibited normal phenotype except a few plants which grew slower than the controls. This may be due to the fact that Fusarium *ftf1* gene is a zinc finger transcription factor. The zinc



finger proteins are present throughout the higher organisms; the siRNAs derived from the partial *ftf1* sequence have influenced the expression of some of the zinc finger genes in banana plant. Moreover, zinc finger proteins are involved in a multitude of physiological processes in plants; it is extremely difficult to pinpoint the cause of this deviation (Ghag *et al.*, 2014).


For pathogenicity studies, both transformed and untransformed plants were inoculated with Foc race 1 at a spore concentration of  $10^6$  mL<sup>-1</sup>. To evaluate the performance of transgenic Rasthali plants under Foc root challenge bioassay, the PCR positive transgenic and non-transformed control plants were transplanted in polybags (30x15 cm) for long term assessment (Sunisha *et al.*, 2020). The plants were monitored for external symptoms and scored periodically. Control plants exhibited yellowing of the older leaves, wilting of lower leaves and pseudostem splitting at 45 days of post inoculation, whereas, no such symptom was observed in transgenic plants (Figure 4a). Expression of dsRNAs targeted against important fungal genes in transgenic plants has been tested successfully in barley, wheat, banana and lettuce (Nowara *et al.*, 2010, Panwar *et al.*, 2013, Ghag *et al.*, 2014 and Govindarajulu *et al.*, 2015). The accumulation of double-stranded or antisense RNA targeting vital fungal transcripts (effector gene *Avra10*) in transgenic barley and wheat has been demonstrated to inhibit the development of the powdery mildew fungus *Blumeria graminis* (Nowara *et al.*, 2010). HIGS for wheat leaf rust fungus caused by *Puccinia triticina* was achieved by targeting three pathogenicity genes of this fungus viz., MAPK, cyclophilin and calcineurin regulatory subunit (Panwar *et al.*, 2013). Similarly, HIGS of HAM34 (Highly Abundant Message #34), CES1 (Cellulose Synthase 1) genes against downy mildew in lettuce was reported by Govindarajulu *et al.* (2015)

Subsequently, after three months of Foc inoculation, 304 transgenic plants exhibited complete wilting (disease severity scale between 2-3) followed by initial yellowing of lower leaves in 84 plants. However, no external Foc symptoms were observed in 46 plants (Figure 4b; Table 2). Both the susceptible and resistant transgenic plants were further subjected to PCR assay for the presence of Foc using specific primer (to amplify SIX gene) and it was found that all the plants had fungal infection (Figure 4 c & d). This indicated that the transgenic plants did not exhibit symptoms despite the presence of fungus in the plants. The RNAi mediated silencing of vital fungal genes (*ftf1*, velvet, ERG6/11) against


Fusarium wilt in banana (Ghag *et al.*, 2014; Dou *et al.*, 2020) supports the findings of this study. The results of this study showed that HIGS can potentially play a major role in controlling Fusarium wilt disease (race 1) in banana, where there is neither a natural resistance gene pool known nor the conventional breeding methods available (Ghag *et al.*, 2012). Moreover, the pathogen control conferred by HIGS is likely to be more durable than resistance mediated by major genes for resistance (Govindarajulu *et al.*, 2015). These 46 *ftf1* transgenic plants were further observed up to 6 months post-inoculation for any delayed Foc infection. Nine plants exhibited complete wilting, followed by marginal yellowing of older leaves in 20 plants. Seventeen transgenic lines were free of Foc-related symptoms even.

**Table 2. Disease severity scale of transgenic plants against Foc race 1 at 3 months post inoculation**


Treatment	No. of plants showing *			
	Grade 0	Grade 1	Grade 2	Grade 3
Rasthali control (Un-inoculated)	30	-	-	-
Rasthali control (Inoculated with Foc)	-	-	1	124
Grand naine control (Un-inoculated)	12	-	-	-
Grand naine control (Inoculated with Foc)	40	-	-	-
Rasthali <i>ftf1</i> (Un-inoculated)	7	-	-	-
Rasthali <i>ftf1</i> (Inoculated with Foc)	46	84	184	120




(a)



(b)



(c)



(d)

**Figure 4.** Bioassay of transgenic banana plants against Foc race 1. a) Infected transgenic plants at 45 days post inoculation, UC 1- wild type Rasthali, UC 2- wild type Grand naine; b) PCR analysis of infected transgenic plants 45 days post inoculation; c) Infected transgenic plants 3 months post inoculation, d) PCR analysis of infected transgenic plants 3 months post inoculation (UC - Un-transformed control).

## CONCLUSION

The results of this study have demonstrated the effectiveness of RNAi technology in imparting resistance to Foc race 1 in banana cv. Rasthali. Further, the durability of the resistance needs to be studied in these disease-free plants. This strategy could potentially be employed in other banana cultivars that are susceptible to Foc race 1.

## Acknowledgment

The authors would like to thank DBT-NER BANANA for the funding and Centre for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University, Coimbatore for providing facilities. RDS also thank DBT-NER banana for the fellowship.

## REFERENCES

- Aurore, G., Parfait, B. and L. Fahrsmann. 2009. Bananas-raw materials for making processed food products. *Trends Food Sci. Technol.* **20**: 78-91.
- Bakry, F., Carreel, F., Jenny, C. and J. P. Horry. 2009. Genetic improvement of banana. In: Jain SM, Priyadarshan PM (eds) Breeding plantation tree crops: tropical species. *Springer Science & Business Media*, Berlin, pp 3-51.
- Bancroft, J. 1876. Report of the board appointed to enquire into the cause of disease affecting livestock and plants. Queensland, 1876. *Votes Proc.* **3**, 1011-1038.
- Baulcombe, D. 2004. RNA silencing in plants. *Nature*, **431**: 356-363.
- Bernstein, E., Caudy, A.A., Hammond, S.M. and G. J. Hannon. 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature*, **409**: 363-366
- Buddenhagen, I. W. 2009. Understanding strain diversity in *Fusarium oxysporum* f. sp. *cubense* and history of introduction of "Tropical Race 4" to better manage banana production. *Acta Hort.* **828**: 193-204.
- Chittarath, K., Mostert, D., Crew, K. S., Viljoen, A., Kong, G., Molina, A. B. and J.E. Thomas. 2018. First report of *Fusarium oxysporum* f. sp. *cubense* tropical race 4 (VCG 01213/16) associated with Cavendish bananas in Laos. *Plant Disease*, **102**: 449.
- Czislowski, E., Fraser-Smith, S., Zander, M., O'Neill, W. T., Meldrum, R. A., TranNguyen, L. T. T., Batley J. and E. A. B.Aitken. 2018. Investigation of the diversity of effector genes in the banana pathogen, *Fusarium oxysporum* f. sp. *cubense*, reveals evidence of horizontal gene transfer. *Mol. Plant Pathol.* **19**: 1155-1171.
- Dale, J., James, A., Paul, J. Y., Khanna, H., Smith, M., Peraza-Echeverria, S., Garcia-Bastidas, F., Kema, G., Waterhouse, P., Mengersen, K. and R. Harding. 2017. Transgenic Cavendish bananas with resistance to Fusarium wilt tropical race 4. *Nat. Commun.*, **14**; **8**(1): 1496.
- Dellaporta, S. L., Wood, J. and J. B. Hicks. 1983. A plant DNA mini preparation: version II. *Plant Molecular Biology Rep.*, **1**: 19-21.



- De Vega-Bartol, J. J., Martín-Dominguez, R., Ramos, B., García-Sánchez, M.-A., and J. M. Díaz-Minguez. 2011. New virulence groups in *Fusarium oxysporum* f. sp. *phaseoli*: The expression of the gene coding for the transcription factor *fff1* correlates with virulence. *Phytopathology* **101**: 470-479.
- Dita, M. A., C. Waalwijk, I. W. Buddenhagen, M. T. Souza Jr. and G. H. J. Kema. 2010. A molecular diagnostic for tropical race 4 of the banana fusarium wilt pathogen. *Plant Pathology*, **59**(2): 348-357.
- Dou, T., Shao, X., Hu, C., Liu, S., Sheng, O., Bi, F., Deng, G., Ding, L., Li, C., Dong, T., Gao, H., He, W., Peng, X., Zhang, S., Huo, H., Yang, Q. and G. Yi. 2020. Host-induced gene silencing of Foc TR4 ERG6/11 genes exhibits superior resistance to Fusarium wilt of banana. *Plant Biotechnol. J.*, **18** (1): 11-13.
- Ghag, S. B., Shekhawat, U. K. S. And T. R. Ganapathi. 2012. *Petunia* floral defensins with unique prodomains as novel candidates for development of *Fusarium* wilt resistance in transgenic banana plants. *PLoS ONE J*, **7**(6): e39557.
- Ghag, S. B., Shekhawat, U. K., & Ganapathi, T. R. 2014. Host-induced post-transcriptional hairpin RNA-mediated gene silencing of vital fungal genes confers efficient resistance against Fusarium wilt in banana. *Plant Biotechnol. J.*, **12**, 541-553.
- Ghildiyal, M. and P. D. Zamore. 2009. Small silencing RNAs: an expanding universe. *Nat. Rev. Genet.* **10**: 94-108.
- Govindarajulu, M., Epstein, L., Wroblewski, T., and R. W. Michelmore. 2015. Host-induced gene silencing inhibits the biotrophic pathogen causing downy mildew of lettuce. *Plant Biotechnol. J.* **13**: 875-883
- Huang, G.Z., Allen, R., Davis, E.L., Baum, T.J. and R. S. Hussey. 2006. Engineering broad root-knot resistance in transgenic plants by RNAi silencing of a conserved and essential root-knot nematode parasitism gene. *Proc. Natl Acad. Sci. USA*, **103**: 14302-14306.
- Jose J. de Vega-Bartol, Raul Martín-Dominguez, Brisa Ramos, Maria-Asuncion Garcia-Sanchez, and Jose Maria Diaz-Minguez. 2011. New Virulence Groups in *Fusarium oxysporum* f. sp. *phaseoli*: The Expression of the Gene Coding for the Transcription Factor *fff1* Correlates with Virulence. *Phytopathology*, Vol. **101** (4): 470-9.
- Mao, Y.B., Cai, W.J., Wang, J.W., Hong, G.J., Tao, X.Y., Wang, L.J., Huang, Y.P. and X.Y. Chen. 2007. Silencing a cotton bollworm P450 monooxygenase gene by plant-mediated RNAi impairs larval tolerance of gossypol. *Nat. Biotechnol.* **25**: 1307-1313.
- Mat Jalaluddin, N. S., Othman, R. Y., and J. A. Harikrishna. 2019. Global trends in research and commercialization of exogenous and endogenous RNAi technologies for crops. *Crit. Rev. Biotechnol.* **39**: 67-78.
- Mustaffa, M. M. and R. Thangavelu. 2010. First report on the occurrence of a virulent strain of Fusarium wilt pathogen (Race- 1) infecting Cavendish (AAA) group of bananas in India. *Plant Disease*, **94**: 1379.
- Muwonge, A., Tripathi, J., Kunert, K. and L. Tripathi. 2016. Expressing stacked HRAP and PFLP genes in transgenic banana has no synergistic effect on resistance to *Xanthomonas* wilt disease. South+A1443 *African Journal of Botany*, **104**: 125-133.
- Nowara, D., Gay, A., Lacomme, C., Shaw, J., Ridout, C., Douchkov, D., Hensel, G., KumLehn, J. and Schweizer, P. 2010. HIGS: host-induced gene silencing in the obligate biotrophic fungal pathogen *Blumeria graminis*. *Plant Cell*, **22**, 3130-3141.
- Nunes, C. C. and Dean, R.A. 2012. Host-induced gene silencing: a tool for understanding fungal host interaction and for developing novel disease control strategies. *Mol Plant Pathol.* **13**(5): 519-529.
- Panwar, V., McCallum, B. and Bakkeren, G. 2013. Host-induced gene silencing of wheat leaf rust fungus *Puccinia triticina* pathogenicity genes mediated by the Barley stripe mosaic virus. *Plant Mol. Biol.* **81**: 595-608.
- Perez-Vicente, L., M. Dita and E. Martinez. 2014. Inoculation of *Fusarium oxysporum* f. sp. *cubeense* causal agent of fusarium wilt in banana. Technical manual prevention and diagnostic of Fusarium wilt (Panama disease) of banana caused by *Fusarium oxysporum* f. sp. *cubeense* Tropical Race 455-58.
- Qi, T., Guo, J., Peng, H., Liu, P., Kang, Z. and Guo, J. 2019. Host-Induced Gene Silencing: A Powerful Strategy to Control Diseases of Wheat and Barley. *Int. J Mol Sci.*, **20**(1): 206.
- Shekhawat, U.K.S., Ganapathi, T.R. and Hadapad, A.B. 2012. Transgenic banana plants expressing small interfering RNAs targeted against viral replication initiation gene display high-level resistance to banana bunchy top virus infection. *J. Gen. Virol.* **93**: 1804-1813.
- Singh, N., Mukherjee, S.K. and M. V. Rajam. 2020. Silencing of the Ornithine Decarboxylase Gene of *Fusarium oxysporum* f. sp. *lycopersici* by Host-Induced RNAi Confers Resistance to Fusarium Wilt in Tomato. *Plant Mol Biol Rep.* **38**: 419-429.
- Strosse H., R. Domergue, B. Panis, J.V. Escalant and F. Côte. 2003. Banana and plantain embryogenic cell suspensions (A. Vézina and C. Picq, eds). INIBAP Technical Guidelines 8. The International Network for the Improvement of Banana and Plantain, Montpellier, France. INIBAP ISBN: 2-910810-63-1.
- Sunisha C, Sowmya H D, Usharani T R, Umeshma M, Gopalkrishna H R, Saxena A. 2020. Deployment of Stacked Antimicrobial Genes in Banana for Stable Tolerance Against *Fusarium oxysporum* f.sp. *cubeense* Through Genetic Transformation. *Mol Biotechnol.*, **62**(1): 8-17.
- Thaler, J.S., Owen, B. and Higgins, V.J. 2004. The role of the jasmonate response in plant susceptibility to diverse pathogens with a range of lifestyles. *Plant Physiol.* **135**: 530-538.

- Thangavelu R, Sundaraju P, Sathiamoorthy S, Raguchander T, Velazhahan R, Nakkeeran S, Palaniswami A. 2001. Status of Fusarium wilt of banana in India. In: Molina AB, Nikmasdek NH, Liew KW (eds) Banana Fusarium wilt management towards sustainable cultivation. INIBAP-ASPNET, Los Banos, Laguna, Philippines, pp. 58–63
- Thangavelu, R., Saraswathi, M.S., Uma, S. 2021. Identification of sources resistant to a virulent Fusarium wilt strain (VCG 0124) infecting Cavendish bananas. *Sci Rep* **11**: 3183.
- Wang, M., Weiberg, A., Lin, F.-M., Thomma, B. P. H. J., Huang, H.-D., and Jin, H. 2016. Bidirectional cross-kingdom RNAi and fungal uptake of external RNAs