



Assessment of Salinity Tolerance in Different Genotypes of Pearl Millet (*Pennisetum glaucum* (L.) R.Br.)

Chilukoti Mallikarjuna¹, Nandhini K^{1*}, Sri Veda Patibandla¹ and Shanthi P²

¹ Agricultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore

² Department of Genetics and Plant Breeding, Tamil Nadu Agricultural University, Coimbatore

Abstract:

This study assessed salinity tolerance in 11 pearl millet genotypes at germination and seedling stages under varying salt concentrations. Results showed a reduction in germination percentage, plumule and radicle lengths, with more significant decreases at higher salinity levels (75mM, 100mM). Despite poor germination, dry weight increased with salt concentration, indicating salt accumulation. Radicle length increased up to 25mM, suggesting plant adaptability to abiotic stresses. Under control conditions, CO 10 and CO (Cu) 9 exhibited the highest germination, while PT 6476 showed the least at 100mM salinity. ICMB 10444 had the maximum plumule length, and ICMB 98222 had the minimum. CO Cu 9 showed the highest radicle length under control, while Kizhilkuppam had the highest at 100 mM salinity. ICMB 15088 had the highest dry weight under control, and CO H 10 had the highest at 100 mM salinity. All genotypes were affected by NaCl stress, reducing seed germination, seedling growth, and nutrient uptake. COH10 showed the least impact, indicating higher genetic potential for salinity tolerance.

Keywords : *Pearl millet genotypes; Salinity tolerance; Germination; Seedling stage; NaCl stress; Dry weight*

INTRODUCTION:

Salinity poses a severe threat to crop production, especially in dry and semi-arid regions worldwide. Soluble salts in soils elevate soil salinity, restricting plant growth by binding water more firmly than plants can extract. Different crops exhibit varying responses to salinity, with most being sensitive, leading to slower growth rates, reduced tillering, branching, and reproductive growth over several months.

India faces substantial salinity challenges, affecting over 6.7 million hectares, including saline, alkali, and coastal soils. Gujarat and Uttar Pradesh bear a significant burden, impacting more than 50% of their lands. The Indo-Gangetic plain, Bihar, West Bengal, Delhi, Haryana, Gujarat, Rajasthan, Madhya Pradesh, Maharashtra, and peninsular states face ecological concerns due to salt-affected soils.

Millets, known as Nutri-Cereals, emerge as vital crops offering abundant nutrition and health benefits. Pearl millet, a standout, contains substantial iron (4-8 mg/100g), aiding in managing anemia. With twice the protein of milk, millets, especially ragi (finger millet), boast the highest



calcium content (364 mg/100g), three times more than milk. Rich in dietary fiber, millets promote digestion, prevent constipation, and possess pre-biotic properties beneficial for gut health.

India is the leading producers of millets Madhya Pradesh, Tamil Nadu, Andhra Pradesh, and Uttarakhand being top-producing states. Millets play a crucial role in India's agricultural landscape.

Salinity Management and Biomass for Renewable Energy:

In dry and semi-arid tropics, managing salinity through soil reclamation or improved irrigation is expensive for underdeveloped nations. Crop improvement emerges as a cost-effective, long-term alternative. Pearl millet, a staple in dry regions, shows promise in saline areas, supporting increased production.

As global energy consumption rises, the adverse environmental effects of non-renewable fuels become evident. Biomass stands out as the renewable energy solution, with millets contributing to biofuel production. Pearl millet, with its superior chemical composition, presents an excellent option for biofuel production due to its high cellulose and hemicelluloses content.

Objective of the Study:

Given the significance of pearl millet, there is a pressing need to expand cultivation into problem soils like saline areas. This study aims to identify salt-tolerant genotypes, screen them under varying saline concentrations, and pinpoint tolerant genotypes during the germination phase. The findings hold the potential to enhance cultivation in saline-prone areas and guide the development of salt-tolerant pearl millet varieties.

Study Objectives:

The study aims to identify salinity-tolerant pearl millet genotypes for increased cultivation in saline areas. Specific goals include genotype collection, screening under different saline concentrations, and identifying tolerant genotypes during germination.

Materials & Methods :

Eleven pearl millet varieties (PT 6476, CO 10, CO H 10, 86 M 38, CO CU 9, KIZHILIKUPPAM, PT 6067, PT 6710, ICMB 10444, ICMB 15088, ICMB 98222) were sourced from the Millets Research Centre, TNAU, Coimbatore.

Characteristics of Pearl Millet Varieties:

- **CO 10:** Tall, strong tillering, quick growth, matures in 70-80 days, 230-250 cm height, high grain production, resistant to downy mildew and shoot flies.
- **CO(Cu)9:** Tall and erect, fast growth, matures in 70-80 days, 250-280 cm height, high grain output, resistant to downy mildew and shoot flies.
- **ICMB 15088:** Early maturity in 70-80 days, variable resistance to pests, medium-sized grains suitable for human and animal consumption.

Salinity Conditions:

Artificial salinity was induced using NaCl at concentrations of 0mM, 25mM, 50mM, 75mM, and 100mM.



Screening at Germination Phase:

NaCl was used at different concentrations (0mM, 25mM, 50mM, 75mM, 100mM) to create salinity conditions. The experiment, conducted in Petri dishes at the Department of Plant Breeding and Genetics, used 11 pearl millet genotypes, with each treatment replicated twice in a factorial randomized block design. Observations recorded eight days after sowing included germination percentage, plumule and radicle lengths, plumule-radicle ratio, and fresh and dry weights.

Observations Recorded (8 Days After Sowing):

- Germination (%)
- Plumule Length (cm)
- Radicle Length (cm)
- Plumule-Radicle Ratio
- Fresh Weight (gm)
- Dry Weight (gm)

Germination Percentage Calculation:

$$\text{GP} = \frac{\text{Seeds germinated}}{\text{Total seeds}} \times 100$$

Statistical Analysis:

Two-way ANOVA was employed to analyse the data, with factors A and B representing levels of interest for the experiment. The model is a fixed-effects model, and the resulting ANOVA table includes factors A, B, their interaction (AB), and error (SSE).

Results and Discussion:

The study focused on evaluating the salt tolerance of 11 pearl millet varieties during the seedling stage. Salinity conditions significantly impacted seed germination, seedling growth, and nutrient uptake across all varieties. COH10 exhibited the least susceptibility, followed by CO(Cu)9, CO10, and ICMB 98222 at the highest salinity level (100 mM), indicating their higher genetic potential for salinity tolerance. Conversely, Kizhilikuppam, ICMB 15088, and ICMB 10444 were identified as highly susceptible genotypes.



The observed reduction in seed germination, seedling growth, and dry matter yields under NaCl stress aligns with previous findings in sorghum and pearl millet. The differential responses among genotypes highlight the importance of genetic factors in salt tolerance. Notably, the increase in dry matter weight with salinity suggests ion accumulation, with COH10 demonstrating resistance to ion toxicity. The study underscores the need for further field research to validate and implement the identified salt-tolerant genotypes in saline-prone areas for developing resilient pearl millet varieties.

CONCLUSION

NaCl stress reduced seed germination, seedling growth, dry matter yields and nutrients uptake by all the eleven pearl millet varieties, but COH10 was least affected compare to other varieties followed by CO(Cu)9 and CO10 and ICMB 98222 under the highest salinity level (100 mM), thus demonstrating its higher genetic potential for salinity tolerance. The genotypes Kizhilikuppam, ICMB 15088 and ICMB 10444 are highly susceptible genotypes. However, further research should be conducted under field conditions in order to authenticate these findings. The tolerant genotypes can be recommended in the saline susceptible area and can be utilized for further development of salt tolerant varieties



Figure .1 Mean percentage of germination for pearl millet genotypes at different levelsof salinity level







Table 2: Mean plumule length (cm) for pearl millet genotypes at different salinity level







Figure 3 Mean Radicle length (cm) for pearl millet genotypes at different levels of salinity level









Figure 4 Mean Dry Weight (gm) for pearl millet genotypes at different levels of salinity level



Seedling evaluation of CO H 10 pearl millet genotype at different levels of salt concentration




Seedling evaluation of CO 10 pearl millet genotype at different levels of salt concentration



Seedling evaluation of Kizhilkuppam pearl millet genotype at different levels of salt concentration



Seedling evaluation of CO Cu 9 pearl millet genotype at different levels of salt concentration





REFERENCES

- Abel.G. H., and A. J. Mackenzie. 1964. Salt tolerance of soybean varieties during germination and later growth. *Crop Sci.* 4, 157- 160.
- Abdul Majid & Poornima Priyadarshini CG .2020. Millet derived bioactive peptides: A review on their functional properties and health benefits, *Critical Reviews in Food Science and Nutrition*, 60:19, 3342-3351
- Akbar, M., and T. Yabuno. 1974. Breeding for saline resistant varieties of rice. II. Comparative performance of some rice varieties to salinity during early developmental stages. *Japan J. Breed.* 24. 176-181.
- Arora, S., and Sharma, V. (2017). Reclamation and management of salt-affected soils for safeguarding agricultural productivity. *J. Safe Agri.* 1, 1–10.
- Ashraf, M., T. Mcneilly, and A. D. Bradshaw. 1986. The potential for evolution of salt (NaCl) tolerance in seven grass species. *New Phytol.* 103, 299- 309
- Chanwala, J., Satpati, S., Dixit, A. et al. Genome-wide identification and expression analysis of WRKY transcription factors in pearl millet (*Pennisetum glaucum*) under dehydration and salinity stress. *BMC Genomics* 21, 231 .2020.
- Emmanuel Iwala, Victor Odjegba, Caroline Umebese, vinay Sharma, Afroz Alam, Physiological and gene expression studies of selected *Zea mays* L. and *Pennisetum glaucum* (L.) R. Br. genotypes to simulated drought stress condition, *Vegetos*, 10.1007/42535-019-00030-7, 32, 3, (397-406), (2019).
- Harshraj Shinde, Ambika Dudhate, Daisuke Tsugama, Shashi K. Gupta, Shenkui Liu, Tetsuo Takano, Pearl millet stress-responsive NAC transcription factor PgNAC21 enhances salinity stress tolerance in Arabidopsis, *Plant Physiology and Biochemistry*, 10. 1016/j.plaphy.2018.11.004. 2018.



- Hussain, K., A. Majeed, Kh. Nawaz, M. Farrukh Nisar, F. Khan, S. Afghan and K. Ali. 2010. Comparative study for salt stress among seed, root stock and direct regenerated violet (*Viola odorata* L.) seedlings in relation to growth, ion contents and enzyme activities. *Afr. J. Biotechnol.*, 9(14): 2108-2117.
- Jamil, M, S. Rehman and E.S. Rha. 2007. Salinity effect on plant growth, PSII photochemistry and chlorophyll content in sugar beet (*Beta vulgaris* L.) and cabbage (*Brassica oleracea* capitatal.). *Pak. J. Bot.*, 39(3): 753-760.
- Jha, R., Mishra, P. Dietary fiber in poultry nutrition and their effects on nutrient utilization, performance, gut health, and on the environment: a review. *J Animal Sci Biotechnol* **12**, 51. 2021.
- Kumar, S., Kumari, R., Singh, V. K., & Rajput, S. K. 2021. Salinity tolerance in pearl millet: current status and future prospects. In *Salinity Stress in Plants* (pp. 147-168). Springer, Singapore.
- Krishnamurthy, L., Serraj, R., Rai, K.N. et al. Identification of pearl millet [*Pennisetum glaucum* (L.) R. Br.] lines tolerant to soil salinity. *Euphytica* 158, 179-188 .2007.
- Mazhar, T., Q. Ali and M.S.R.A. Malik. 2020. Effects of salt and drought stress on growth traits of *Zea mays* seedlings. *Life Sci. J.*, 17: 48-54.
- O. P. Yadav, K. N. Rai, Genetic Improvement of Pearl Millet in India, *Agricultural Research*, 10.1007/40003-013-0089-z, 2, 4, (275-292). 2013.
- Roshan Kumar Singh, Mehanathan Muthamilarasan, Manoj Prasad, Biotechnological approaches to dissect climate-resilient traits in millets and their application in crop improvement, *Journal of Biotechnology*, Volume 327,2021, Pages 64-73, ISSN 0168-1656,



R.K. Maiti, Luz Elena Maiti, Sonia Maiti, Angelica Maria Maiti, M. Maiti, Humberto Maiti, Genotypic Variability in Maize Cultivars *Zea mays* L.) for Resistance to Drought and Salinity at the Seedling Stage, *Journal of Plant Physiology*, Volume 148, Issue 6, 1996, Pages 741-744, ISSN 0176-1617

Shweta Jha, Plant Functional Genomics Lab, Biotechnology Unit, Department of Botany (UGC-Centre of Advance Study), Jai Narain Vyas University, Jodhpur 342001, India

Siddique, M. A., Jaiswal, D. K., & Srivastava, S. 2020. Physiological and molecular responses to salinity stress in pearl millet. *Journal of Plant Biochemistry and Biotechnology*, 29(4), 599-611.

Singh, G., & Kumar, A. (2019). Breeding pearl millet for salt tolerance: present status and future prospects. *Indian Journal of Genetics and Plant Breeding*, 79(4), 692-702.

Yakubu, H., A.L. Ngala and I.Y. Dugje. 2010. Screening of millet (*Pennisetum glaucum* L.) varieties for salt tolerance in semi-arid soil of Northern Nigeria. *World J. Agric. Sci.*, 6(4): 374-380.

Yakubu, H., H. Usman and A.M. Chiroma. 2006. Screening of sorghum varieties for salt tolerance in a sandy loam soil of Borno State, Nigeria. *Niger. J. Exp. Appl. Biol.*, 7(1):49-54