



Estimation of Nutritive Composition of *Sauropus androgynous* (Multi Vitamin Plant) at Different Growth Stages and Position of Leaves

***Naveena E ¹ , Janavi G J¹, Rameshkumar D², Senthilnathan S³ , and Savitha B K⁴**

1 Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore -641003.

2 SRM College of Agricultural Sciences, SRM Institute of Science Technology, Vendhar Nagar, Baburayanpettai, Chengalpattu-603201

3 Department of Agricultural Economics , Agricultural college and Research Institute, Tamil Nadu Agricultural University , Killikulam, Vallandu, Tamil Nadu, India

4 Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore -641003.

***Corresponding Author: naveenaep@gmail.com**

Abstract

A research was designed to estimate the nutritive composition of Chekkurmanis (Katuk) leaves on two different growth stages viz., 60 and 120 days after planting and two different positions viz., terminal and basal whorls in plants. Utilization of green leafy vegetables differs from leaves of annuals, shrubs and trees. Katuk leaves rich in vitamins and minerals and hence known as protective food. The results of the study showed that Vitamin C, Protein, Calcium, Carbohydrate contents increased with the growth stages from 60 to 120 days after planting. In contrast, Vitamin A content decreased as the plant approached 120 days. There is a significant difference in nutritive value from 60 to 120 days after planting. Vitamin A was observed to be significantly high (4.11 mg/100g) in terminal whorl at 60 days after planting compared to in the terminal whorl at 120 days (3.01 mg/100g). Higher Vitamin A (5.12 mg/100g) was observed in basal whorl leaves at 60 days after



Madras Agric.J., 2023; <https://doi.org/10.29321/MAJ.10.200D15>

(first draft)

planting, while Vitamin C (220.41 mg/100g), Protein (4.99 g/100g), Calcium (4.16%), Carbohydrate (9.83 g/100g) were observed in basal whorls at 120 days after planting. In conclusion, the basal whorls had more embedded nutritive value than the terminal whorl leaves. This research reveals that nutritive composition of *Sauropus androgynus* mainly depends on their growth stages and leaf position in the plant.

Keywords: Chekkurmanis leaves, Nutritive analysis, Growth Stages, Position (terminal or basal whorl) of leaves

Introduction

Chekkurmanis (*Sauropus androgynus* L.), belongs to the family Euphorbiaceae, it has vernacular names viz., *Katuk* or *Sweet leaf bush*, it is a shrubby glabrous perennial green leafy vegetable, extensively grown in warm humid tropics with ample rainfall. The crop is native of Indoburma region, later introduced to India from Malaya. It is widely distributed in Sikkim, Himalaya, Khasi, Abour and Arka hills, Western Ghats of Kerala. The plant is also cultivated in Australia, as it is nutritious, most prolific, high yielding with appetizing green leaves. Katuk is mainly propagated through semi hardwood cuttings of 20 - 30 cm length stem. The tender leaves and succulent shoots are used for cooking. In India, it is commonly known as Multivitamin or Multi green plant, because of its rich nutritive value, containing large amount of essential mineral elements, with high level of Vitamin A, Vitamin B, Thiamine (B1), Riboflavin (B2), Vitamin C, Carbohydrate, Calcium, Potassium, Phosphorous, Iron and Dietary Protein (Jeevitha *et al.*, 2022). Due to presence of antioxidants in the leaves, is utilized for treating many diseases viz., diabetics, cancer, allergy, microbial infection, cholesterol and swelling as reported by Pant *et al.* (2020). In India, anciently Katuk leaves were used to improve the eyesight, to cure skin diseases, urinary problems, cardiovascular problems, relieves internal fever and several other illness as stated by Ong (2003). Mineral deficiency and other related diseases can be overcome by intake of leafy vegetables at recommended quantity, since an appropriate amount of vitamins, minerals and phytochemicals are necessary for normal functioning of human metabolic processes. To understand the nutritive composition of Chekkurmanis at different growth stages and position of leaves, a research was carried out at the Horticultural College and Research Institute, Periyakulam.



Materials and methods

The healthy and disease-free rooted cuttings of *Sauropus androgynus* collected from Orchard, Agricultural College and Research Institute, Tamil Nadu Agricultural University (TNAU), Madurai, Tamil Nadu, were used as planting materials for the research work. Field trial was taken up at Horticultural College and Research Institute, Periyakulam. The experiment was conducted at Field No. 26 of Western Block Farm, Department of Vegetable Science, Horticultural College and Research Institute, Periyakulam, which is located at 80° East longitude, 11° North latitude, at an altitude of 300 M above Mean Sea Level. The nature of soil in the experimental site was sandy loam. Chekkurmanis leaf samples from terminal and basal whorls of the plants were collected at 60 and 120 days after planting, to assess the nutritive composition of young (terminal whorls) and matured (basal whorls) leaves. The leaves were washed with tap water to remove adhering sand, dust and other contaminants on the surface. Fresh leaf samples were used for analyzing vitamins, carbohydrate and protein, while the leaf samples dried in hot air oven at 60° C for six hours were used for calcium analysis. All the analyses were carried out in three replication. Estimation of Vitamin A was done based on the protocol developed by Jensen (1978). Similarly, Estimation of Vitamin C described by Harris and Ray (1935), Estimation of Carbohydrate by anthrone method suggested by Hodge and Hofreiter (1962), Estimation of Protein were analyzed separately by Biuret method as per Layne (1957) and Estimation of Calcium analysis by versenate method suggested by Jackson (1973).

Statistical analysis

All the analyses were performed in triplicate and Analysis of variance (ANOVA) was carried out using statistical package WINDOWSTAT to evaluate variation between two levels of growth stage and leaf position of the plant.

Result and discussion

Vitamin A

The fresh leaves from terminal and basal whorl of *Sauropus androgynus* contained 4.11mg/100g and 5.12mg/100g Vitamin A respectively on 60 DAP, while 3.01mg/100g and 4.26mg/100g on 120 days after planting. There existed significant



(first draft)

difference ($P < 0.05$) in Vitamin A content in terminal and basal whorl leaves on 60 and 120 days after planting. High Vitamin A content was found in basal whorl leaves on 60 days after planting compared to terminal leaves on 120 days after planting. This trend of Vitamin A content in Chekkurmanis leaves is analogous with results reported by Padmavathi and Rao (1990) and less than the values reported by Devi *et al.* (2007). It was found that vitamin A content was considerably reduced with later growth stage of the plant. This ensures the fact that Vitamin A is redistributed to every organ of plant as reported by Hocmuth *et al.* (2004). It is essential for effective functioning of visual system, reproduction, immune system, growth and development as stated by Sommer and WHO (1995).

In the present research work, Vitamin A content of Chekkurmanis decreased as the plants advanced its growth. Similar report was noticed in *C. argentea* by Adegbaaju *et al.* (2019) and Biesiada *et al.* (2007) for Leek, Zucchini and Kohlrabi. The values obtained from this research reveals that the higher Vitamin A was found than *Amaranthus digitata*, *Hibiscus sabdariffa* and *Vigna unguiculata* as noticed by Patricia *et al.* (2014), Sathya *et al.* (2021) and moringa leaves by Abbas *et al.* (2018).

Vitamin C

The Vitamin C content at terminal and basal whorl of fresh Chekkurmanis leaves were 112.26 mg/100g and 123.33 mg/100g respectively on 60 DAP, while on 120 days after planting the content were 199.30 mg/100g and 220.41 mg/100g respectively. The Vitamin C content increased considerably with age. There existed significant difference ($P < 0.05$) in Vitamin C content in the terminal and basal whorl leaves on 60 and 120 days after planting. High Vitamin C (220.41 mg/100g) content was noted in basal whorl leaves on 120 days and lesser value (92.26 mg/100g) in terminal leaves on 60 days after planting. Similar trend was observed in *Telifairia occidentalis* as reported by Patel and Srinivasan (2017), Musa *et al.* (2012) and Jeevitha *et al.* (2022). The previous study in Chekkurmanis by Padmavathi and Rao (1990) and Singh *et al.* (2011) reported that vitamin C content was 244.00 mg/100g and 314.30 mg/100g respectively in terminal and basal whorls, which was higher than this research. According to Lee and Kadar (2000), the maturity of leaves is a major factor that determines the compositional value of vegetables. Further Korus (2010) observed that Vitamin C content depends on degree of plant maturity and variety. In this research, Vitamin C attained the maximum value of 220.41 mg/100g in basal whorl



(first draft)

leaves on 120 days. It is very high when compared with *Amaranthus digitata* (70.00 mg/100g) and *Hibiscus sabdariffa* (30.00 mg/100g) as reported by Patricia *et al.* (2010) and *Moringa oleifera* (0.56 mg/100ml) as reported by Achikanu *et al.* (2013).

Vitamin C supports synthesis of hormone, iron absorption and collagen, which is extensively used in cosmetics and pharmaceuticals. It acts as water soluble antioxidant, which plays a major role in free radical scavenging activity and activates immune system (Locato *et al.*, 2013).

Table 1. Nutritive composition of terminal and basal leaves of Chekkurmanis leaves

| Days after planting (DAP) | Position of leaf | Vitamin A (mg/100g) | Vitamin C (mg/100g) | Carbohydrate (g/100g) | Protein (g/100g) | Calcium (%) |
|---------------------------|------------------|---------------------|---------------------|-----------------------|-------------------|-------------------|
| 60 DAP | Terminal whorl | 4.11 ^b | 112.26 ^b | 5.71 ^b | 2.63 ^b | 1.69 ^b |
| | Basal whorl | 5.12 ^a | 123.33 ^a | 6.20 ^a | 3.04 ^a | 2.08 ^a |
| 120 DAP | Terminal whorl | 3.01 ^b | 199.30 ^b | 8.81 ^b | 4.25 ^b | 2.94 ^b |
| | Basal whorl | 4.26 ^a | 220.41 ^a | 9.83 ^a | 4.99 ^a | 4.16 ^a |
| SEd | | 0.114 | 4.492 | 0.139 | 0.101 | 0.056 |
| CD (P < 0.05) | | 0.280 | 10.992 | 0.342 | 0.248 | 0.137 |

Mean values carrying superscript letters represent significant difference at P < 0.05

Table 1 represents the nutritive composition of terminal and basal whorls of Chekkurmanis leaves at two different growth stages viz., 60 and 120 days after planting and two positions in plants.

Carbohydrate

The carbohydrate value observed in terminal and basal whorl of fresh leaves were 5.71g/100g and 6.20g/100g respectively on 60 days, while 8.81g/100g and 9.8 g/100g respectively on 120 days after planting. Carbohydrate content was high (9.83g/100g) in basal leaves on 120 days and low (5.71g/100g) in terminal leaves on 60 days. The basal whorl leaves consisted of more carbohydrate (6.20g/100g and 9.8g/100g) when compared with terminal whorl leaves (5.71g/100g and 8.81g/100g) on 60 and 120 days. There existed significant difference (P < 0.05) in carbohydrate content on the terminal and basal whorl leaves on 60 and 120 days. The content also significantly increased with respect to the maturation of the plants and this finding is in line with the findings of Adegbaaju *et al.*



(first draft)

(2019). The carbohydrate quantity observed in this current research was identical with earlier research reported in *Sauropus androgynus* by Chakraborty *et al.* (2019), but lower than as reported by Devi *et al.* (2007). Carbohydrate is a fundamental element for normal functioning of human body. Ingestion of carbohydrate enriched leaves adds energy level to diet by providing favourable nutrition. Amount of carbohydrate in basal leaves of katuk is higher than *T. occidentalis* but lower than *C. aconitifolius* as indicated by Otitoju *et al.* (2014) and equal to the *Senna occidentalis*, *Solanum nodiflorum*, *Physalis viscosa* by Odhav *et al.* (2007) and in moringa leaves as observed by Abbas *et al.* (2018).

Protein

Fresh leaves at terminal and basal whorl leaves of Chekkurmanis contained protein content in the range of 2.63 g/100g and 3.04 g/100g respectively at 60 days, while it was 4.25 g/100g and 4.99 g/100g respectively at 120 days after planting. There existed significant difference ($P < 0.05$) in protein content on the terminal and basal whorl leaves on 60 and 120 D A P. Previously, Singh *et al.* (2011) published the literature and reported that protein content of Chekkurmanis leaves were 5.25 g/100g. From the result of this study, the protein content of *Sauropus androgynus* increased with age of the plant. According to Bamishayie (2011), the protein content of moringa leaves was moderately low at initial stage (27.61%) and finally reached higher values on later stage (28.08%). The highest protein content (4.99 g/100g) was recorded in basal whorl of the plant on 120 days after planting and the lowest content (2.63 g/100g) was recorded on terminal whorl of the plant 60 days after planting. The protein values found in basal leaves of Chekkurmanis on 120 days after planting were higher than *Petroselinum crispum* (2.97 g/100g), *Anethum graveolens* (3.46 g/100g), *Lactuca sativa* (1.62 g/100g), *Brassica oleraceae* (1.21 g/100g) as reported by Caunii *et al.* (2010), but lower than *Moringa oleifera* leaves (6.7 g/100g) as reported by Gopala Krishnan *et al.* (2016). Therefore, *Sauropus androgynus* has the potential to serve as a very good source of protein for the reduction of protein malnutrition. It also acts as neurotransmitter and carries the oxygen in blood. Deficiency of protein causes malnutrition diseases viz., kwashiorkor and marasmus and results in stunted growth. It is also useful for immune function, creation of enzymes and hormones and maintains fluid balance in the body.



Calcium

Calcium content of dried leaf samples collected from both terminal and basal whorl of leaves was in the range of 1.69% and 2.08% respectively on 60 days, whereas it was 2.94% and 4.16% respectively on 120 days. There existed significant difference ($P < 0.05$) in calcium content in the terminal and basal whorl leaves on 60th and 120 days after planting. Calcium percentage obtained in this study is more (4.16%) than that (0.7%) as stated by Padmavathi and Rao (1990). In the present study, the highest calcium content observed in basal leaves at 120 days was 4.16%. This finding is in agreement with Modi (2007), who observed more calcium content in long day interval than short day interval in amaranthus plant. In contrast, the lowest calcium content of 1.69% was recorded in the terminal leaves on 60 days. The highest calcium content in basal leaves might be due to immobility (non-translocation) of calcium in the plant. This concept is in agreement with the authors Taiz and Zeiger (2002) and Hochmuth *et al.* (2004).

When the plants approach maturity, the nutrient content (calcium) in the leaves gets increased. Similar trend is in agreement with the values reported by Platel and Srinivasan (2017) for Chekkurmanis crop and similar tendency with age of plant was found in tender and mature stages of *Celosia argentea* plant by Adediran *et al.* (2015). Calcium plays an important role in nervous system, cell signaling and muscle contraction. It also helps to maintain healthy teeth and bones. Calcium is also utilized by cells for transferring ions and triggers several enzymes secretion in human body by Sadler (2011). The values found in this study are higher than some of underutilized green leafy vegetables such as *Digera arvensis* (0.50%), *Amaranthus tricolor* (0.23%) and *Cucurbita maxima* (0.30%) as reported by Gupta *et al.* (2004), *Hibiscus sabdariffa* (0.4%) and *Ipomea aquatica* (0.09 %) by Singh *et al.* (2011), *Alternanthera sessilis* (2.0 %) and *Celosia argentea* (2.03%) as reported by Rekha Sinha (2018) and *Moringa oleifera* (1.14%) by Srivastava *et al.* (2018).

Conclusion

The results of the research indicated that *Sauropus androgynus* is an excellent source of Vitamin A and C, Carbohydrate, Protein and Calcium than any other leafy vegetables. Further, the green leafy vegetables are fine source of vitamins, minerals and other minor elements. The research reveals that basal whorl of leaves consist of more



Madras Agric.J., 2023; <https://doi.org/10.29321/MAJ.10.200D15>

(first draft)

nutrient content than terminal whorl leaves and there existed significant difference between the nutritional constituents in leaves upon growth of the plant.

Funding and acknowledgment

The authors are grateful to the Department of Vegetable Science, HC & RI, Periyakulam for providing technical support of our research

Ethics statement

No specific permits were required for the described field studies because no human or animal subjects were involved in this research

Consent for publication

All the authors agreed to publish the content

Competing interests

There is no conflict of interest in the publication of the content

Author's contribution

Research grant – GJG, EN, Idea conceptualization- GJG, Experiment – EN, writing original draft- EN, DRK, Data analysis- DRK, reviewing and editing – SS, BKS.

References

- Abbas RK, Elsharbasy FS, Fadlelmula AA. 2018. Nutritional Values of (*Moringa oleifera*) Total Protein, Amino Acid, Vitamins, Minerals, Carbohydrates, Total Fat and Crude Fiber, under the Semi-Arid Conditions of Sudan. *Journal of Microbial and Biochemical Technology.*, **10(2)**: 56-58. DOI: 10.4172/1948-5948.1000396.
- Achikanu CE, Eze-Steven PE, Ude CM, Ugwuokolie OC. 2013. Determination of the vitamin and mineral composition of common leafy vegetables in South Eastern Nigeria. *International Journal of Current Microbiol. Appl. Sci.*, **2(11)**: 347-353. DOI: 10.4236/jss.2016.43010
- Adediran OA, Gana Z, Oladiran JA, Ibrahim H. 2015. Effect of age at harvest and leaf position on the yield and nutritional composition of (*Celosia argentea* L.) *International Journal of Plant and Soil Science.*, **5(6)**: 359-365. DOI:10.9734/IJPSS/2015/15063
- Adeggbaju OD, Otunola GA, Afolayan AJ. 2019. Proximate, mineral, vitamin and anti-nutrient content of (*Celosia argentea*) at three stages of maturity. *South African Journal of Botany.*, **24**: 372-379. DOI:10.1016/j.sajb.2019.05.036



Madras Agric.J., 2023; <https://doi.org/10.29321/MAJ.10.200D15>

(first draft)

- Amanabo Musa, Emmanuel Ogbadoyi O. 2012. Effect of Plant Leaf Positions on Some Micronutrients, Anti-nutrients and Toxic Substances in (*Telfairia occidentalis*) at the vegetative phase. *American Journal of Experimental Agriculture*; **2(2)**: 219 - 232.
- Bamishaiye EI, Olayemi FF, Awagu EF, Bamshaiye OM. 2011. Proximate and phytochemical composition of (*Moringa oleifera*) leaves at three stages of maturation. *Advance Journal of Food Science and Technology*., **3(4)**: 233-237. DOI: 10.4236/fns.2015.611103.
- Biesiada A, Kołota E, Adamczewska-sowinska K. 2007. The effect of maturity stage on nutritional value of leek, zucchini and kohlrabi. *Veg. Crops Res.*, **66**: 39–45.
- Caunii A, Cuciureanu R, Zakar AM, Tonea E, Giuchici C. 2010 Chemical composition of common leafy vegetables. *Studia Universitatis" VasileGoldis" Arad. Seria Stiintele Vietii. Life Sciences Series.*, **20(2)**: 45-48. DOI:10.1016/j.pbi.2007.04.009
- Chakraborty Ivi , Riadh Ilahy, Balaji Vikram, Arghya Mani. 2019 Recent Trends & Advances in Food Science& Post Harvest Technology Satish serial publishing house, Delhi., **1(1)**: 207-218. Doi : 10.1016/B978-0-12-800872-0.00018-4
- Devi NS, Sadhan Kumar PG, Peter KV, Indira V. 2007. Indigenous leaf vegetables for administering vitamin A and minerals. In International Conference on Indigenous Vegetables and Legumes. Prospectus for Fighting Poverty, Hunger and Malnutrition. *Acta Hort.*, **752**, 367-372. DOI: 10.17660/ActaHortic.2007.752.64
- Gopalakrishnan L, Doriya K, Santhosh Kumar D. 2016. *Moringa oleifera*: A review on nutritive importance and its medicinal application. *Food science and human wellness.*, **5 (2)**: 49-56. Doi: 10.1016/j.fshw.2016.04.001
- Gupta S, Jyothi Lakshmi A, Manjunath MN, Prakash J. 2005. Analysis of nutrient and antinutrient content of underutilized green leafy vegetables. *LWT-Food Science and Technology.*, **38(4)**: 339-345. Doi: 10.1016/j.lwt.2004.06.012
- Harris LJ, Ray SN. 1935. Determination of plasma ascorbic acid by 2, 6-dichlorophenol indophenol titration. *Lancet.*, **1(7)**: 1. doi: 10.1042/bj0290275
- Hocmuth G, Maynard D, Vavrina C, Hanlon E, Simonne E. 2004. Plant tissue analysis and interpretation for vegetable crops in Florida. University of Florida's Institute of Food and Agricultural Sciences (UF/IFAS)., **1**:79. Doi: 10.32473/edis-ep081-2004
- Hodge JE, Hofreiter BT. 1962. In: Methods in Carbohydrate Chemistry, Academic Press, New York. DOI: 10.4236/ns.2019.113009



Madras Agric.J., 2023; <https://doi.org/10.29321/MAJ.10.200D15>

(first draft)

Jackson ML. 1973. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd. New Delhi., 183-204. DOI: 10.4236/me.2018.96073

Jeevitha, S, G. J. Janavi, J. Prem Joshua, V. Vani, and T. Anitha. 2022. "Influence of drying methods on nutritional composition of Chekurmanis (*Sauropus androgynus* L.).

Jensen A. 1978. Chlorophylls and carotenoids. Handbook of phytological Methods, Cambridge Univ. Press, London., 59- 70. DOI: 10.4236/fns.2015.63030

Kalpana Platel, Krishnapura Srinivasan. 2017. Nutritional Profile of Chekurmanis (*Sauropus androgynus*), A Less Explored Green Leafy Vegetable. *Indian Journal of Nutrition and Dietetics.*, **54 (3)**: 244. DOI:10.21048/ijnd.2017.54.3.15765

Korus A. 2010, Level of vitamin C, polyphenols and antioxidant and enzymatic Activity in three varieties of kale (*Brassica oleracea* L. var. *acephala*) at different stages of maturity. *Int. J. Food Properties.*, **14**: 1069-1080. DOI:10.1080/10942910903580926

Layne E. 1957. Methods Enzymol. Academic Press, New York -3, 447-466. DOI: 10.4236/ajac.2015.67058

Lee SK, Kader AA. 2000. Preharvest and postharvest factors influencing vitamin C content of horticultural crops. *Postharvest Biol. Technol.*, **20**: 207-220. DOI:10.1016/S0925-5214(00)00133-2

Locato V, Cimini S, DeGara L. 2013. Strategies to increase vitamin C in plants: from plant defense perspective to food biofortification. *Front. Plant Sci.*, **4 (152)**: 1-12. DOI:10.3389/fpls.2013.00152

Modi AT. 2007. Growth temperature and plant age influence on nutritional quality of Amaranthus leaves and seed germination capacity. *Water SA.*, **33(3)**: 369-376. DOI:10.4314/wsa.v33i3.180598

Odhav B, Beekrum S, Akula US, Baijnath H. 2007 Preliminary assessment of nutritional value of traditional leafy vegetables in KwaZulu-Natal, South Africa. *Journal of Food Composition and Analysis.*, **20(5)**: 430-435. DOI:10.1016/j.jfca.2006.04.015

Ong HC. Sayuran: Khasiat Makanan & Ubatan, Utusan.2003. Publications and Distributors, Kuala Lumpur, Malaysia. Doi: 10.1155/2015/714158

Otitoju GTO, Ene-Obong HN, Otitoju O. 2014. Macro and micro nutrient composition of some indigenous green leafy vegetables in south-east zone Nigeria. *J Food Process Technol.*, **5(11)**: 1-6. DOI:10.4172/2157-7110.1000389



Madras Agric.J., 2023; <https://doi.org/10.29321/MAJ.10.200D15>

(first draft)

Padmavathi P, Prabhakara Rao M. 1990. Nutritive value of *Sauropus androgynus* leaves.

Plant Foods Hum Nutr., **40(2)**: 107-13. DOI:10.21048/ijnd.2017.54.3.15765

Patricia O, Zoue L, Megnanou RM, Doue R, Niamke S. 2014. Proximate composition and nutritive value of leafy vegetables consumed in Northern Cote d'Ivoire. *European Scientific Journal.*, **10 (6)**: 212-227. DOI: 10.19044/ESJ.2014.V10N6P

Rekha Sinha. 2018. Nutritional analysis of few selected wild edible leafy vegetables of tribal of Jharkhand, India. *Int. J. Curr. Microbiol. App. Sci.*, **7(2)**: 1323-1329. DOI: 10.20546/ijcmas.2018.702.161

Sadler TW. 2011. Langman's Medical Embryology. Wolters Kluwer Lippincott Williams & Wilkins, Edn 11, Philadelphia. Doi: 10.1016/j.tplants.2010.07.003.

Sathya M, Sumathi P, Senthil N, Vellaikumar S, John Joel A. 2017. Direction of association among β -carotene, grain yield and its component traits in RIL population of pearl millet. 2017. *Electronic Journal of Plant Breeding*, **8 (2)**: 548-553. DOI: 10.5958/0975-928X.2017.00082.5

Singh S, Singh DR, Salim KM, Srivastava A, Singh LB, Srivastava RC. 2011. Estimation of proximate composition, micronutrients and phytochemical compounds in traditional vegetables from Andaman and Nicobar Islands. *International Journal of Food Sciences and Nutrition.*, **62(7)**: 765–773. DOI:10.3109/09637486.2011.585961

Sommer A, 1995. World Health Organization. Vitamin a Deficiency and its Consequences: A Field Guide to Detection and Control. World Health Organization, Edn 3, Geneva.

Srivastava, Pan RS, Bhatt BP. 2018. Antioxidant and nutritional potential of some underutilized leafy vegetables consumed by tribals of Jharkhand, India. *Current Science.*, **114(6)**: 1222-1233. DOI:10.18520/cs/v114/i06/1222-1233

Taiz L, Zeiger E. 2002. Plant physiology. Sinauer Associates, Inc. Sunderland. Massachusetts. Edn., **3**: 370-372. Doi: 10.3389/fnut.2022.861347

Usha Pant, Ram Bhajan, Anita Singh, Kalpana Kulshethra, Singh A K, Himanshu Punetha. 2020. Green leafy mustard: A healthy alternative. *Electronic Journal of Plant Breeding*, **11 (01)**: 267-270. DOI: 10.37992/2020.1101.045.