

R. C. Wood. I wish you all success. It seems a long time to my Coimbatore days—after all it is a good many years, and I am getting nearer the time when I am due to retire, but I have very pleasant recollections of the time I spent there.

Telegraphic Messages were received from: Rao Bahadur B. V. Nath, P. H. Rama Reddy, Principal Lyalpur Agricultural College and others.

Messages were received from: Mr. K. Gopalakrishna Raju; The Vice Chancellor, Andhra University; Director of Agriculture Hyderabad State; Director of Agriculture, Bombay; and Mr. Kasiviswanatha Thampuram.

CULTURAL METHODS OF CONTROLLING PLANT DISEASES

BY RUDOLPH D. ANSTEAD, M.A., C. I. E.,

Retired Director of Agriculture, Madras Presidency.

Since the day, sixty years ago, when the Madras Agricultural Department was born, views on many agricultural problems have undergone a profound change. This is especially the case in those branches of the subject which deal with the pests and diseases of crops. During the past fifty years enormous strides have been made in medical knowledge of all kinds, including the health of plants.

An incessant war is carried on between man and insects, fungi, and bacteria, and many are the methods which have been recommended to combat these pests which take an enormous annual toll of our crops and stored products, and also of life of man and beast.

Despite this agricultural practices have been remarkably little influenced. It seems so obviously the right thing to ascertain the nature and life history of a pest and then to attack its weakest and most vulnerable phase. This, however, does not get to the real root of the problem, and in most cases is only a palliative. The hosts of the enemy remain, undiminished at their source, and the remedies have to be constantly applied. It is now being realised that direct attack by assault and battery is nearly always useless, and entomologists and mycologists are being rapidly transformed into plant pathologists, bringing these subjects into line with new developments of medical thought. A more insidious technique has begun to appear, which may be called perhaps the "cultural" method of preserving plants in health. The presence of the pest is ignored in this technique, and no direct attack is made on it.

In his Presidential Address to the Agricultural Section of the British Association at Toronto as long ago as 1924 Sir John Russell said: "these cultural methods of dealing with plant diseases and pests offer great possibilities, and the close study jointly by plant physiologists and pathologists of the response of the plant to its surroundings, and the relationships between the physiological conditions of the plant and the attack of the various parasites would

undoubtedly yield results of great value for the control of plant diseases."

Mycologists and Entomologists are turning their attention more and more to the effects of soil and climate on the incidence of disease, and it is now becoming generally recognised that there are vast possibilities of controlling many plant diseases, not by attacking the disease organisms themselves, but by controlling conditions in such a way that these organisms are unable to develop because they find the conditions imposed deleterious to them. Though the organisms are present they are unable to become effective because the conditions are not favourable to them.

For example, MacRae pointed out that foot rot (*Helminthosporium*) of Wheat in Northern India occurs only on early sown fields, and the remedy is not to spray, but to delay sowing until the cold weather sets in, and the temperature imposes conditions unfavourable to the development of the fungus. This is a purely "cultural" remedy based on a study of conditions which favour the crop and are unfavourable to the disease.

It is well known that a plant, or an animal, in good vigorous health is resistant to disease attack when subjected to infection. It is the plants or animals which are ill nourished and weak which fall easy victims. Hence the plant pathologist has in recent years turned his attention more and more to the study of the factors which maintain a plant or animal in vigour. When these are known it is often possible to provide for resistance when an epidemic of some sort, insect or fungoid, come along.

In 1924-25 the demonstration areas under cotton at Chendathur in the Fourth Circle were perfectly healthy, while all round the cotton on the ryots fields was attacked by "black arm" and looked as if a fire had been through them. All that had been done on our demonstration areas was to employ correct cultural methods. Dr. C. L. Withycombe dealing with the "frog hopper" pest of Sugarcane in the West Indies said that, "canes do not necessarily show serious" blight when foghoppers have been abundant, nor is an abundance of "the insect a necessary condition for serious blight", and he maintained that the controlling factor was often the presence of plenty of water physiologically available to the canes, a factor which could be arranged for. Again, cotton leaf-spot (*Altenaria longipediciliata*) is a weak parasite able to infect weak tissue only under the most favourable circumstances, and yet in Trinidad when cotton is water-logged or has poor root growth it becomes a serious pest. (Empire Cotton Growing, Review. V. 1.48.)

Tunstall when reviewing Tea diseases and their remedies (Quar. Jour. Indian Tea Association 1920) puts cultural methods, such as

improved drainage, removal of excessive shade, and clean pruning, before direct methods like spraying, and work in Ceylon has shown that tea bushes which fail to recover after pruning and are attacked by *Diplodia* are really deficient in reserves of food. Wallace again, concludes that all the available evidence points to "leaf scorch", a frequent cause of loss to orchard growers, arises from defective nutrition and unsatisfactory water supply, cultural defects which can be remedied by drainage and manuring. (Jour. Pomology & Hort. Science VII 1 & 2.)

Rotation of crops will sometimes prevent disease attack. A case in point is the Betel Vine in the Madras Presidency which when grown continuously on the same land is apt to become infested with *Phytophthora* wilt disease, absent when rotation is practiced.

Eelworm attack on Sugar Beet and Potatoes is a danger. In Germany in 1876 this pest became so widespread that twenty-four sugar beet factories had to be closed down. The remedy lies in rotation of crop. Beet should only be grown once in four years on the same land.

Another method is that adopted by the plant breeder who, in many cases has been able to evolve new strains highly resistant to particular diseases so that the actual presence of insects or micro-organisms may be ignored. One of the latest examples of this method is the evolution of a "blast" (*Piricularia*) resistant strain of paddy at Coimbatore. Many other examples could be quoted. The strains of wheat resistant to rust produced at Cambridge by Sir R. H. Biffin are world famous, whilst varieties of potatoes resistant to virus diseases, and Poplar hybrids resistant to canker are well known.

An interesting example in this direction is the case of apple scab. At one time it was thought that this fungus pest could only be controlled by constant spraying, but experiments at the East Malling Research Station in Kent (England) have shown that certain rootstocks induce resistance, while others induce susceptibility to the disease. Hence it is possible to select rootstocks on which to graft apples which will help the grower to ward off the scab disease by a cultural method, and spraying is then unnecessary, or at any rate more effective. It is of interest to note that trees which were well manured benefitted from spraying more than trees on starved land. On the latter the disease is apt to be so bad that any control is impossible.

The internal condition of the food plant in relation to insect attack is of importance. The association of particular species of insects with particular food plants has resulted in an adaptation on the part of the insect with regard to the physiology of its digestion in a manner best suited to its requirements. Many insects fail to live on other than their normal food plants. The resistance or immunity of a plant to

insect attack is often due to factors closely associated with the physiology of the plant, probably the presence or absence of particular substances in the tissues of the plant. Thus Andrews showed that the vitality of *Helopeltis*, the "mosquito blight" of Tea is directly controlled by the suitability or otherwise of the food supply, and when a constant supply of soluble potash is applied to the roots of the tea bush it will remain immune from attacks for a long time.

Sugar Beet develops a specific disease in the absence of Boron: Oats suffer from a grey fleck disease in the absence of Manganese, though only one part in a million may be necessary to prevent this: Zinc appears to be essential for fruit trees which are otherwise attacked by rosette disease.

This leads to the question of vitamins which have been found to be so essential to the health of animals and man. Pioneer work carried out by the Madras Agricultural Department by Lt. Col. McCarrison, Viswanath, and others has indicated that there is a relationship between the supply of vitamins and the organic content of the soil, and has emphasised the importance of maintaining the humus content of soil. (Mem. Dept. of Agri. in India. Chem. Series. IX. 27. Indian Jour. Medical Research. IV. 4.)

The plant apparently obtains vitamins from the organic matter, possibly directly, and these vitamins are handed on to the animals which feed on them. The author would suggest that it is within the bounds of possibility that the vitamins are just as important to the health of the plant as they are to the health of the animals, and that it is not likely that the plant is merely acting as a transferring medium for these essentials of health. There is a growing mass of evidence to prove that when the humus content of the soil is allowed to run down below a certain level crops become increasingly subject to diseases of all kinds. Hence the importance of the use of organic fertilisers like Activated Composts.

Sir Albert Howard claims that in another fifty years time all plant diseases will be dealt with along such lines as have been here indicated, and that spraying machines and the like will only be found in museums. Though the author is not prepared to go quite so far as that, he does maintain that in the future more and more attention will be devoted in the campaign against plant pests and diseases to the cultural method of attack rather than to the shock attack of the sprayer, and he trusts that the Madras Agricultural Department, which he had the honour and privilege at one time to serve, will be found at the end of the next fifty years in the forefront of the battle in the same proud position which it has occupied since the day it was founded.