

SELECTED ARTICLE

Development of Modern Composting Methods.

By Y. D. Wad, Institute of Plant Industry, Indore, Central India.

The deliberate use of rotted organic wastes—vegetable and animal—for the purpose of growing larger quantities of better crops seems to be as old as the art of agriculture itself. It is conceivable that primitive man may have simultaneously (1) noticed the superiority of natural vegetation growing in virgin forest land rich in organic matter, and (2) discovered the possibility of artificial cultivation of some of the plant species suitable for his food.

At any rate, the importance of farmyard manure to crop growth has been stressed in ancient Indian and European literature on agriculture (Russell and Richards, 1917). Kind (1926) has described in detail how the Chinese peasants of old, took elaborate care to collect all available wastes and convert them systematically into well-rotted composts. It is noteworthy that in every part of the world this system of returning its own waste material to land has maintained soil fertility in spite of continuous cropping through the ages. The crowded population of China is still being maintained on the produce of its soil after its agricultural use for over forty centuries. This is perhaps the most convincing proof of the perfect balance of ancient systems of agriculture with their environment. It is very striking, indeed, that modern composting technique has very little to add to the basic principles underlying the Chinese method of making manure from agricultural wastes.

Liebig published in 1840 his essay *Chemistry in its application to Agriculture and Physiology*. This marked the beginning of a period when scientific investigations and commercial enterprise concentrated on the stimulation of crop production by means of factory-made chemical manure. Subsequent work at Rothamstead and elsewhere established the manufacture of artificial fertilizers on a sound footing. Factories engaged during the war in the fixation of atmospheric nitrogen needed new markets afterwards. This further intensified the use of chemical manures. The use of bulky farm manures fell into the background. It was even asserted that this practice was not an essential feature of agriculture. A school of scientific workers, however, soon arose who maintained that a certain proportion of humus is essential to preserve the crumb structure in soils and that such a structure in turn was essential for efficient plant growth (Russell, 1934; *Symposium on Soil Organic Matter*, 1927).

Another group of scientists (Howard, 1937, 2) believed that the artificial stimulation of soil activities for commercial cropping was sure to upset the natural balance of soil factors and in the long run might lead to evils not yet fully realized. They maintain, therefore, that in any agricultural system adequate provision is absolutely necessary for returning all the waste products of agriculture back to the land. Howard (1937, 1) even maintains that in specialized systems such as the planting industries it may be necessary to make provision for the supply of humus to the soil by manufacturing it at extra cost from other sources to enable the soil to meet the abnormal strain resulting from highly intensive cultural practices.

Also, the large majority of the cultivators in the world still believe that the produce obtained by the use of chemical manures is not always equal in quality to that obtained by the use of ordinary farm manure.

Recent discoveries of workers on animal nutrition have apparently confirmed this belief by their findings (McCarrison, 1926, 1937; Viswanath and Suryanarayana, 1927; Ramiah, 1933). It has also been claimed (Howard, 1937,1) that the use of humic manures from vegetable and animal wastes imparts disease resistance both to crops and the animals that feed on them. Recently, a fresh impetus was received by the investigations into the nature of soil humus and the decomposition of organic wastes to humus (Russell and Richards, 1917; Waksman et al., 1929; Du Toit and Page, 1930, 1932; Waksman and Iyer, 1932, 1933; and others). This was accompanied by zealous attempts of other workers to discover how to make larger quantities of humic manures and how to increase the speed of the decomposition (Carbery and Finlow, 1928; Rao and Subrahmanyam, 1932, 1935; Anstead, 1932; Gadgil and Hegdekatti, 1937). These workers aimed at ensuring a copious supply of cheap and properly made humic manure.

Richards and Hutchinson (1921) artificially converted straw to humus by the help of ammonium sulphate. This led to the development of the patented "Adco" process.

Fowler (1930) and Howard concentrated their efforts on the utilization of all available organic residues for making composts of the Chinese type. Fowler stressed that it is necessary to build up an intensively active biological starter of the proper type to ensure a good start and maintain the speed thus secured throughout the course of decomposition. His system of making "activated composts" is founded on this principle and is applicable equally to both farm residues and town wastes.

Howard aimed chiefly at making all types of residues into composts and thus increasing the supply of cheap humus. He saw in this a means to compensate for the existing shortage of cattle dung for manure-making in India where cattle dung is badly needed for fuel purposes in the absence of a satisfactory substitute. His work in this direction culminated the development of Howard and Wad's Indore Process (1931, 1935).

This process aims at utilizing the harder residues by making them less refractory to the influence of the fermenting microorganisms by the physical cracking of tissues or by exposing them to the corrosive environment of actively decaying material of a better composition. It lays special stress on starting the heap with a physical structure capable of maintaining adequate aeration without undue loss of moisture all through the period of decay notwithstanding its compaction due to the shrinkage of the rotting mass. It is maintained that a properly made heap will very soon develop within it all the required intensity of microbiological activity by itself. All the temperature ranges and sequences of the types of micro-organisms necessary for composting will automatically appear. The process is aerobic, clean and sanitary as well as cheap and simple. The final product always maintains the proper standard of quality.

This process, therefore, spread rapidly all over the world and is applicable to a large variety of cultural systems and environments. It can convert all types of wastes quickly into well-rotted composts. This is typically illustrated by its application (1) to the disposal of habitation wastes (Jackson and Wad, 1934; Howard, 1935, 1937, 1938), (2) the composting of cane trash (Tambe and Wad, 1935; Dymond, 1923, 1938) and of sisal wastes, the wastes of tea, coffee, rubber and coconut and oil palms (Bagot, 1937; Howard, 1938) and its modifications for making composts with rain water (Timson, 1939) and by the intermittent supply of water from canals (Jackson, Wad, and Paise, 1934).

Fowler (1930) seems to have considered partially anaerobic conditions during the later stages of decomposing heaps as having some beneficial effect.

The author of this note has observed that under the hot arid climate of the Rajputana desert the compost made with three turns had an inferior chemical composition than that produced by one turn only. It appeared that due to the different degrees in the course of fermentation of the various components of the heap the more easily decaying portions under the stimulus of local climate reached the stage of complete oxidation and consequent losses by the time the more refractory parts were sufficiently crumbled down. It is possible that losses of this nature may be kept down by lessening the number of turns or altering their intervals to regulate the ventilation to the desired degree.

While investigating the possibilities of the hot fermentation process Rajgopal *et al.* (1936) have concluded that in compost heaps a better conservation of carbon and nitrogen is possible when anaerobic conditions follow after a vigorous aerobic start with rise of temperature. The mechanisms by which this is brought about is yet to be fully investigated.

Howard (1937) has recently evolved what he calls "Sheet Composting". This seems to suit wherever labour is scarce or costly. Residues of field crops are composted *in situ* in the field without collecting and removing them. The conditions in sheet composting are perhaps semi-aerobic. The following description by Howard will illustrate the application of this principle:—

This development was worked out during the last two years on the potato areas of South Lincolnshire which have been noted to suffer from shortage of humus. After the pea-crop grown for canning has been harvested, the land is immediately drilled with beans. The sown area is then covered with a layer of crushed straw from the shelling machines followed by a thin layer of farmyard manure. The Indore process then sets in on the surface of the soil. The beans grow through the fermenting mass and at the end of September are ploughed in with the layer of finished compost. Decay is rapid and by the time the fields are planted in potatoes the following spring the resulting humus has been incorporated in the soil and is ready for nitrification. This modification is known as sheet composting—the making of humus in a thin layer all over the surface. Catch crops of beans or mustard or a crop of weeds can also be manured with humus or farmyard manure before ploughing in in the autumn when sheet composting again takes place. The turf of old pastures or old leys can be converted into humus in a similar fashion. The Indore process has in this way been applied with success to no less than three important practical problems, green manuring, the effective utilization of weeds and stubble and the better utilization of the old turf of grass land.

Similar attempts at simplification are being made by applying waste organic matter direct to the soil with inorganic reinforcements (Eden, 1935, 1936). The present system of burying tea prunings and loppings of shade trees along with the chemical manures may also be considered a similar operation.

It appears to the author that perhaps the most economic and convenient method of returning waste material to land will be a preliminary aerobic decomposition to suitable stage followed by direct application to the field, a few weeks before sowing time, before preparatory cultivation begins. There seems to be some scope for such a method as it involves the least deviation from current routine as well as the minimum of labour and care. *Agriculture and Livestock in India* Vol. IX Part V.