Some Additions to the Tingid Fauna of South India

The lace wing bugs belonging to the family Tingidae (Hemiptera) have been studied extensively elsewhere. A preliminary list of the Tingid fauna of South India was published in 1962 (Mohanasundaram, 1962). then, several species have been either collected or identified from previous collections. The present note is based on the collections of lace wing bugs made in South India by the author during the period 1962-70 and from the unidentified tingid specimens retrieved from the mixed insect collections of the Entomology Section, Agricultural College and Research Institute, Coimbatore.

1. Cysteochila delineata (Distant) (Distant, 1910, p. 119) Syn. Bredenbachius delineatus Distant.

The species was described from Calcutta and later collected from Bauhinia purpuria.

Specimens are available as collected from a red flowered tree from Bangalore by T. B. Fletcher on 26-8-1913.

2. Bako malayanus (Drake)

Syn. Galeotingis malayana Drake

Recorded on Panicum repens, Cynodon dactylon, Cenchrus glaucus, Oryza sativa and Saccharum officinarum by Mohanasundaram (1962). The present specimens were collected from Bamboo leaves in Coimbatore which is a new host record.

3. Amniams ravanus (Kirkaldy)

Syn. Phyllontochila ravana Kirkaldy (Distant 1904, p. 136)

This species was recorded from Ceylon on Vitex trifolia. Specimens are on hand collected by P. S. Nathan, from Talliparamba, N. Malabar during July-August, 1918. Host has not been given.

4. Monosteria edeia Drake & Livingstone Drake & Livingstone (1964)

This species was described from Zizyphus jujuba in North India.

Several specimens have been collected from the same host from Vizagapatnam (Andhra Pradesh), July 1965.

5. Naochila sufflata (Drake & Poor)

(Drake & Ruhoff, 1965)

This was originally recorded on Ehretia laevis and Lantana sp. from Dehra Dun area.

Specimens were collected from a tree belonging to *Ficus* sp. from Oilseed Research Station, Salem. September 1964.

Hegesidemus otiosus Drake

(Drake & Ruhoff, 1965; p. 241)

It was recorded from Nilgiri Hills without host record.

The present specimens were collected at Doddapetta Peak in Nilgiris at 8000 feet on *Strobilanthus* plants on 11-9-1970.

7. Paracopium cingalanse (Walker)

Syn. Dictyonota cingalanse (Distant, 1904; p. 128)

This species has been previously recorded from India and Ceylon. E. J. Woodhouse had found it causing hypertrophy of the corolla of *Clerodendron phlomoides* flowers in Bengal.

Specimens are on hand collected by N. K. M. in Coimbatore on 8-7-1938 labelled as "in fruits of Chlorodendron chlomidis". The floral cecidia in this case has been wrongly conceived as fruits

 Tingis (Caenotingis) beesoni Drake (Drake & Ruhoff, 1965; p. 405)

Recorded from Burma on *Gmelia* arborea. Specimens are on hand labelled as "Vizagapatam, Anakapalli, 26-12-1928, V. T. Rao Coll. on *Gmelia* arborea".

ACKNOWLEDGEMENTS

The author wishes to thank the Entomologist and Professor of Entomology, Agricultural College and Research Institute, Coimbatore for permitting to peruse the Section collections and for the encouragement given in the preparation of this note.

M. MOHANASUNDARAN

Agricultural College and Research Institute, Colmbatore 641003.

REFERENCES

- DISTANT, W. L. 1904 Fauna of British India Rhynchota, Vol. II. Taylor and Francis, London.
- DISTANT, W. L. 1910. Fauna of British India.
 Rhynchota. Vol.V. Heteroptera: Appendix.
 Taylor and Francis, London.
- DRAKE, C. J. & DAVID LIVINGSTONE, 1964 The new species of lace bugs from India. (Hemiptera-Tingidae). The Great Basin Naturalist XXIX (1): 27-30.
- DRAKE, C. J. & F. A. RUBOFF. 1965. Lace bugs of the World. A catolog (Hemiptera: Tingidae) Smithsonian Institution, Washington.
- MOHANASUNDARAM, M. 1962. A preliminary list of the Tingid Fauna of South India. (Tingidae: Hemiptera). Madras agric. J. 49 (11): 365 70.