

Characteristics and the yield potential of different pepino accessions

J. MOHAN AND M. SUBASH CHANDRA BOSE

Horticultural Research Station, Tamil Nadu Agri. University, Ooty - 643 001, Tamil Nadu

Abstract : Pepino (*Solanum muricatum*) is a small herbaceous perennial minor fruit cum vegetable crop with multiple possibilities for consumption. Special features include its high yield potential, easy propagation by stem cuttings, perennial nature, fast growing and high response to intensive cultivation and fertilizer application and above all the habit of flowering and fruit set round the year. As on date 9 promising accessions have been established based on plant and fruit morphology, yield performance and quality traits. These accessions also offered differing preference for different utilities such as dessert fruit, squash etc. (**Key Words :** Pepino, *Solanum muricatum*, Fruit morphology, Solanaceae, Minor fruit)

Pepino (*Solanum muricatum*), often referred as Melon Pear, is a small herbaceous perennial belonging to Solanaceae. It is sub-tropical, medium to large sized minor fruit species native to Northern Andes, South America. For the first time in India it was introduced at Horticultural Research Station, Ooty during the latter parts of 1994.

Pepino is a nutritious vegetable cum fruit crop with multiple possibilities for consumption like green and cooked vegetable, vegetable salads, pachadies, fresh fruit (dessert), fruit salad, delicious fruit juice, squash, etc. Pepino fruits are highly juicy and moderately sweet and possesses characteristic pleasing aroma like that of melon fruit. These fruits are rich in minerals, vitamin C, free amino acids and organic acids. It contained comparatively low starch and low soluble sugars. The other constituents include inositol, lipids, pigments, cellulose, hemicellulose and pectic substances.

Pepino prefers cooler climate (temperature between 15 and 25 C) and performs well in fertile loamy soil with good drainage. It is easily propagated through stem cuttings. It responds well to high level management and it is a high fertilizer responsive crop. Even the quality parameters showed considerable variation in response to soil fertility, manuring and fertilization. According to Heyes et al. (1993) pepino is

a potentially useful research tool for investigation. Pluda et al. (1993) stated that changes in nutrient composition and salinity have marked influence on fruit quality i.e. fruit quality may be improved by crop management.

Materials and Methods

Since its introduction in 1994 at horticultural Research Station, Ooty, pepino was multiplied through both open pollinated seeds and stem cuttings. The plants raised from seeds showed variations in leaf and fruit morphology and based on desirable traits and yield potential, 9 accessions were chosen and multiplied through vegetative propagation. The yield and quality (fruit and juice) evaluations have shown very interesting results and are presented hereunder.

Results and Discussion

The data on leaf characteristics are presented in Table 1. Accessions SMu-4 and 5 produced trifoliolate leaves from the very beginning and the accessions SMu-1, 6, 7, 8 and 9 produced entire leaves in the early stages and tri- or penta-foliolate leaves subsequently. The accessions SMu-2 and 3 continued the production of entire leaves for a longer time and latter produced trifoliolate leaves. The mean leaf length and breadth also showed variations among the accessions; the smallest leaves were observed in SMu-2.

Table 1. Leaf characteristics of pepino accessions

Accession No.	Leaf Morphology	Leaf length (cm)	Leaf breadth (cm)
SMu-1	Entire/Trifoliolate	14.45	3.88
SMu-2	Entire	8.40	2.30
SMu-3	Entire	12.00	3.03
SMu-4	Trifoliolate	12.00	3.13
SMu-5	Trifoliolate	14.13	3.53
SMu-6	Entire/Trifoliolate	10.65	3.50
SMu-7	Entire/Trifoliolate	14.90	4.10
SMu-8	Entire/Trifoliolate	15.40	4.50
SMu-9	Entire/Trifoliolate	13.50	3.67

The fruit morphological characteristics (Table 2) exhibited striking variations. The fruit shape varied from nearly round to long oblong. The accessions SMu-4, 6 and 7 are characterized with muricated surface where as others possessed smooth surface. The characteristic violet or purple streaks were absent in accessions SMu-2, 7, 8 and 9. The colour of young fruits varied from marble white to dark green and that of the ripefruits from light yellow to red. Pluda *et al.* (1993) have stated

that some clones excel in some agronomic and quality characteristics, sharing some common traits such as erect growth habit, heavy fruit set, good colour, size and acceptable flavour. Fruit yield varied from 48.95 t ha⁻¹ in SMu-5 to 68.40 t ha⁻¹ in SMu-1. The accession SMu-1 recorded highest yield followed by SMu-5 indicating the significant variability of fruit yield of different accessions.

Table 2 Fruit characteristics of pepino accessions

Accession No.	Morphology		Colour		Flesh Colour	Mean fruit wt. (g)	Yield (t ha ⁻¹)
	Shape	Presence of purple streaks	Unripe	Ripe			
SMu-1	Oblong	P	Green	Golden yellow	Yellowish red	276	68.40
SMu-2	Round	N	Marble white	Golden Yellow	Red	128	50.30
SMu-3	Long oblong	P	Dark green	Golden Yellow	Light Yellow	152	55.05
SMu-4	Muricated Round	P	Light Yellow Or marble White	Light Yellow	Yellow orange	53	53.70
SMu-5	Round/oblong	P	Light green to Marble white	Red yellow	Reddish yellow	157	48.95
SMu-6	Muricated oblong	P	Green	yellow	yellow	95	58.15
SMu-7	Oblong	N	Green	Yellow	Light yellow	157	58.80
SMu-8	Muricated round	N	Green Reddish	Reddish yellow	145	52.15	
SMu-9	Tomato shaped	N	Green	Yellowish red	Yellow orange	140	51.25

Note : P =Purple streak present; N = No streak

Mean fruit weight of pepino varied from 53 g in SMu-4 to 276 g in SMu-1. Maroto *et al.*, (1997) reported fruit weights varying from 145-300 g. The striking feature is that under Ooty conditions the fruit size in all the 9 accessions are grouped under Class I as they all weigh more than 50 g and have good shape. Ercan and Akili (1996) classified pepino fruits according to their weight and shade into two classes viz., Class I: Fruits with good appearance and each weighing 50 g each. It is

interesting to note that seed set is considerably higher under the Nilgiris climatic conditions. The seed number per fruit varied from few tens to few hundreds. This may be one of the main reasons for the extremely large size of fruits (a maximum weight of 742 g was recorded at Woodhouse farm) Ercan and Akili (1996) reported that hand pollinated plants produced fruits with an average weight of 115.48 containing 8 seeds, compared to control which produced seeds less fruits 73.44 g

mean weight. Pluda *et al.* (1993) observed marked fruit quality variations with change in nutrient composition and levels of salinity. According to Heyes *et al.* (1994) pepino is a potential research tool for investigation.

Among the fruit and juice quality parameters (Table 3) dry matter content, electrical conductivity, sugar content and titrable acidity showed considerable variation among the accessions but fruit specific gravity, juice pH and juice density showed little

variation. Dry matter content of fruits varied from 18.52% (SMu-8) to 27.13% (SMu-5); EC from 2.26 mS/cm² (SMu-4); sugar content from 6.0 to 6.5% (SMu-7 and 8) to 9.0 to 11.0% (SMu-5) and titrable acidity from 12.8 me/100ml (SMu-5) to 18.0 me/100 ml (SMu-4). These results are in accordance with the reports of Redgwell and Turner (1986), who have reported an elaborate account of chemical quality parameters.

Table 3. Fruit and juice quality variations of pepino accessions

Accession No.	Fruit			Juice			
	Specific gravity	Dry matter content (%)	EC (mS/cm)	PH	Juice density (g/cc)	Sugar%	Titrable acidity (me/100ml)
SMu-1	0.983	26.87	2.70	6.00	1.028	8.0-10.	013.6
SMu-2	0.890	25.12	2.78	5.89	1.002	7.0-8.5	14.0
SMu-3	0.914	24.28	2.92	5.70	1.010	7.5-9.0	14.4
SMu-4	0.926	20.12	3.12	5.59	1.019	7.0-7.5	18.0
SMu-5	0.910	27.13	2.81	5.97	1.025	9.0-11.0	12.8
SMu-6	0.928	26.03	2.76	5.89	1.006	8.0-9.0	14.4
SMu-7	0.862	21.16	2.26	5.80	1.014	6.0-6.5	16.8
SMu-8	0.892	18.52	2.75	5.95	1.017	6.0-6.5	16.4
SMu-9	0.906	19.82	2.71	5.94	1.015	6.5-7.2	13.9

The visual quality attributes such as size, shape, symmetry, colour of rind, colour of flesh, olfactory, physical traits like firmness and structure and chemical attributes like flavour, sweetness, acidity etc., are conducive for consumer preference. Accordingly the accessions SMu-1 and 5 offer better preference for dessert purposes because of the higher dry matter and sugar content, lower titrable acidity and better shape and eye appealing colour. SMu-1 and 7 were found to have higher preference for squash and vegetable purposes. Thus among the 9 accessions, SMu-1 which is almost similar to colossal variety of New Zealand, seems to be the most ideal one and the accessions SMu-5 may be preferred for dessert purposes.

References

- Ercan, N and Akili, M. (1996). Reasons for parthenocarphy and the effects of various hormone treatments on fruit set in pepino (*Solanum muricatum* Ait.). *Scientia Hort.* 66 : 141-147.
- Heyes, J.A., F.H. Blaikie, C.G. Downs and D.F. Sealey (1994). Textural and physiological changes during pepino (*Solanum muricatum* Ait.) ripening. *Scientia Hort.* 58 : 1-15.
- Maroto, J.V., Lopez-Galaza, B., Pascual, M.S. Bono, A. San Bautista and J. Alagarda (1997) Procarpil Enhances Earliness and Parthenocarpy of pepin (*Solanum muricatum*) Hort. Sci. 32 : 133.
- Pluda, D., H.D. Rabinowitch and U. Kafkafi, (1993). Pepino dulce (*Solanum muricatum* Ait.) - Quality characteristics respond to nitrogen nutrition and salinity, *J. Ame. Soc.Hort. Sci.* 118 : 86-91
- Redgwell, R.J. and Turner, N.A. (1986) Pepino (*Solanum muricatum*) chemical composition of ripe fruit. *J. Sci. Food Agri.* 37 : 1217-1222.

(Received : July 1999 ; Revised : June 2001)